



Hatching Distribution and Embryo Mortality of Eggs Laid by Broiler Breeders of Different Ages

■ Author(s)

Almeida JG
Vieira SL
Reis RN
Berres J
Barros R
Ferreira AK
Furtado FVF

Departamento de Zootecnia
Universidade Federal do Rio Grande do Sul

■ Mail Address

Sergio Luiz Vieira
Faculdade de Agronomia – UFRGS
Av. Bento Gonçalves, 7712
91.540-000. Porto Alegre, RS, Brazil
Phone/Fax: +55 51 3308 7419

E-mail: slvieira@ufrgs.br

■ Keywords

Broiler chicks, embryo mortality, incubation, post-hatching management.

Projeto com suporte financeiro da FAPERGS.

Arrived: March / 2006
Approved: May / 2008

ABSTRACT

This study investigated the hatching distribution of eggs from broiler breeders of different ages in distinct periods of incubation. Eggs were incubated in a single-stage experimental incubator. A number of 3.510 eggs were distributed into 3 treatments with 13 replicates of 90 eggs each. Treatments corresponded to breeder age: young (34 weeks), intermediate-age (44 weeks) and old (72 weeks) breeders. Eggs were transferred to the hatcher at 432 incubation hours, hatching was first checked at 449 hours, after which the number of hatched chicks was counted every 6 hours up to 515 incubation hours. After each count, hatched chicks were removed from the hatcher. Data were submitted to analysis of variance using measures repeated. A significant interaction between breeder age and incubation time was found. The total period required to hatch all incubated fertile eggs was not influenced by breeder age, which, however, affected hatching distribution. Eggs from old breeders hatched later as compared to young and intermediate-age breeders. More than 71% of the eggs had already hatched at 485 incubation hours, and 94% at 491 hours. Eggs laid by old breeders presented higher infertility and total embryo mortality, resulting in lower hatching percentage.

INTRODUCTION

Commercial hatcheries apply incubation periods of approximately 504 hours, which supposedly maximizes hatching. However, many chicks hatch earlier, and therefore remain for several hours with no food or water before being removed from the hatcher. According to Vieira & Pophal (2000), broiler chicks hatch in a time range of 480 to 510 hours. Factors such as breeder age, hatchable egg storage time and temperature, egg weight, season, and bird type influence total incubation time (Wilson, 1991).

Fanguy *et al.* (1980) and Wyatt *et al.* (1985) showed that the longer the chicks remain in the hatcher, the worse is their live performance. Wyatt *et al.* (1985) observed that chicks that remained in the hatching trays for 14 to 32 hours were 5-32% lighter than those removed 7 hours after hatching. Nir & Levanon (1993) and Sklan *et al.* (2000) found higher percentage of body weight loss, from 5 to 15%, which continued up to market age. Halevy *et al.* (2000) showed that post-hatching fasting lead to lower weight gain and breast meat accretion capacities.

Chicks that hatch and are immediately removed from the hatcher and supplied with water and feed present better performance as compared to chicks that remain in the hatcher longer or take longer to have access to feed and water after hatching (Williams *et al.*, 1951; Kingston, 1979; Fanguy *et al.*, 1980; Hager & Beane, 1983; Wyatt *et al.*, 1985). Gonzales *et al.* (2003) showed that weight loss, early mortality,



lack of flock uniformity, and culling significantly increase after 48 hours of fasting.

The period immediately after hatching is critical for the development of the immune and gastrointestinal systems. During the first days of the bird's life, the small intestine grows five times faster than the rest of the body and small intestine microvilli grow significantly faster in birds supplied with water and feed immediately after hatching (Dibner *et al.*, 1998).

Broiler performance during the first week of life is highly correlated with performance at market age (Nitsan, 1995). Therefore, any factor that reduces initial weight gain can affect body weight at market age.

Wilson (1991) reported that eggs laid by old breeders tend to have long incubation periods, as well as those from breeders at the onset of lay. According to Marques (1994), eggs from younger breeders tend to have a shorter incubation period. Opposite results were reported by Shanawany (1984), who observed that eggs laid by old breeders have shorter incubation periods as compared to those laid by young breeders. That author concluded that from old breeder eggs present earlier embryo development during the first weeks of lay as they remain longer in the oviduct, leading to earlier hatching. These findings are consistent with studies of McNally & Byerly (1936), Smith & Bohren (1975), and Burton & Tullet (1985), who observed shorter incubation periods as breeders aged.

Pedroso *et al.* (2005) observed that eggs laid by 37-week-old breeders hatched 9 hours later than those laid by 32-week-old breeders. In addition, egg weight had no influence on incubation period, which was related to breeder age. These results agree with Vieira *et al.* (2005), who observed that breeder age influences the time required for hatching. Working with breeders of different ages (29, 40, and 59 weeks), those authors found that at 485 hours of incubation more than 90% of the eggs had already hatched. Reis *et al.* (1997) studied 33- and 49-week-old breeders, and verified that 56% of the eggs hatched at 485 hours, and at 490 hours, more than 94% had hatched in a total incubation period of 504 hours. Hudson *et al.* (2004) worked with breeders of different ages (29; 41; 53 e 65 weeks), and observed that 27% of all eggs hatched at 480 hours, and that 86% at 491 hours. According to those authors, the low hatching rate at 480 hours was due to young breeders' (29 weeks) eggs - only 10% had hatched at that time. That study also found that the time required for hatching increased as breeders aged - eggs from 53-week-old breeders or older presented longer incubation periods. This is

consistent with the finding of Olsen (1942); Williams *et al.* (1951), and Wilson (1991), who also verified longer incubation periods as breeders aged. Vieira *et al.* (2005) observed that 90% of eggs hatched at 485 hours remained for 19 hours in the hatcher to allow the remaining 10% to hatch.

Embryo mortality rates must be reduced to improve hatching rates. It is essential to determine its causes and to solve the problems that may occur from laying to hatching.

Breeders with extremes in age usually present high percentage of infertile eggs (Schmidt *et al.* 2003) and high total embryo mortality (McDaniel *et al.*, 1979; Wilson, 1991; Reis *et al.*, 1997; Lapão *et al.*, 1999; Hudson *et al.*, 2004).

The objective of the present study was to determine possible differences in the time required for hatching and embryo mortality of eggs laid by heavy breeders of different ages.

MATERIAL AND METHODS

A total of 3.510 Isa Vedette eggs laid by breeders of three different ages (34, 44, and 72 weeks) were collected in the same farm and were laid on the same day. Eggs were incubated in a single-stage machine in the incubator of the lab of the Animal Science Department of the Federal University of Rio Grande do Sul. Eggs were hourly turned, and submitted to 37.5°C temperature and 60% relative humidity (RH) in the incubator and 36.5°C and 65% RH in the hatcher after, transference on day 18 of incubation. Each breeder age was subdivided in three treatments of 13 replicates each, with 90 eggs per replicate randomly distributed in the machine. The number of hatched chicks started to be counted at 449h of incubation at regular 6-h intervals up to 515h. Two eggs per replicate (n=78) were collected to determine albumen, yolk, and eggshell ratios. Humidity loss during incubation was calculated as the egg weight difference between egg weight at the time of incubation and egg weight time of transference to the hatcher. Egg weight to chick weight ratio was determined by weighing 20% of hatched chicks in each replicate. All birds that hatched up to 485 incubation hours in each replicate were weighed, and 2 were sacrificed; the remaining birds were kept in the hatcher and weighed again at 515 incubation hours, when 2 other birds per replicate were sacrificed to weigh the yolk sac. This allowed the determination of chick weight to yolk sac weight ratio and to verify the effect of 30 h of fasting on these



parameters. Non-hatched eggs from each replicate was analyzed at the end of the experimental period for embryo mortality causes and period, which was divided in early (1-7 days), intermediate (7-14 days), and late (14 a 21 days) mortality.

The study model included "time", as the 6-h intervals when the completely hatched chicks were removed from the hatcher (incubation period). Data to analysis of variance using repeated measures over time in the Mixed Procedures of SAS (1998) software. Variance and covariance structures were determined according to Akaike's evaluation criterion (1987), and compound symmetry (CS) was the one that best fit the data. This structure assumes variance homogeneity for each age. Differences among means were verified by the test of ($p < 0.05$).

RESULTS AND DISCUSSION


Hatching time results as a function of breeder age are shown in Tables and Graphs 1 and 2. The tables show significant interaction between breeder age and hatching time, indicating that the time required for hatching depends on breeder age. In Table 1 and Graph 1, the data are based on the number of hatched chicks, and the hatching percentage at each evaluated time is added (cumulative). Table 1 shows that the first hatchings happened at 461h, from eggs of old and intermediate-age breeders. Young breeders' eggs hatched later, at 467 incubation hours. Eggs from intermediate-age breeders presented higher hatching percentage at 467 incubation hours as compared to those of young and old breeders, which were not significantly different from each other.

After 473 incubation hours, the hatching percentage of young breeders' eggs was statistically similar to those laid by intermediate-age breeders, and remained as such until the end of the experimental period (515 hours). The hatching percentage of old breeders' egg was the lowest up to 485 incubation hours, after which there were no differences.


Difference in hatch distribution between did not affect total incubation period. At 485 incubation hours, 71% of the eggs had hatched. It was observed that eggs from old breeders lowered this average, with only 58% hatching at that time as compared to 77% of young and intermediate-age breeders' eggs.

Based on this observation, it is suggested that if old breeders' eggs are incubated a few hours before the eggs of young and intermediate-age breeders, hatchings could be synchronized at 485 or 491

incubation hours. Therefore, chicks can be removed from the hatcher in two different steps: at 485 or 491 hours and at 515 hours. In the present experiment, 94% of the hatched chicks could be removed at 491 incubation hours, 24 hours before the final incubation time. The remaining 6% could be removed at 515 incubation hours, thereby preserving chick quality.



Graph 1 - Cumulative hatching of chicks derived from broiler breeders of different ages (34, 44 e 72 weeks), % relative to total number of hatched chicks.



Graph 2 - Cumulative hatching of chicks derived from broiler breeders of different ages (34, 44, and 72 weeks), % relative to total number of incubated eggs.

Table 2 and Graph 2 show hatching percentage to the number of incubated eggs, and reflect the results presented in Table 1 and Graph 1. Table 2 shows the differences in hatching percentage at 515 incubation hours according to breeder age. It is observed that young and intermediate-age breeders' eggs presented the best hatching results, and were not statistically different. The worst results were found for old breeders' eggs, which hatching percentage was 18% lower as compared to the eggs from breeders of the other ages.

These results are consistent with those of Pedroso *et al.* (2005) and Vieira *et al.* (2005), who observed a



significant relationship between hatching distribution and breeder age, but disagree with the studies of Reis *et al.* (1997) and Hudson *et al.* (2004), who did not find such relationship.

Table 1 - Cumulative hatchings of chicks derived from broiler breeders of different ages (34, 44, and 72 weeks), % relative to total number of hatched chicks.

| Incubation (hours) | Breeder | | | | Incubation (mean) | CV (%) |
|--------------------|---------------------|--------------------|--------------------|--------------------|---------------------|--------|
| | P | YB | IB | OB | | |
| 449 | 1,0000 | 0,00 | 0,00 | 0,00 | 0,00 | 0,00 |
| 455 | 1,0000 | 0,00 | 0,00 | 0,00 | 0,00 | 0,00 |
| 461 | 0,5207 | 0,00 | 0,21 | 0,23 | 0,15 ^e | 384,06 |
| 467 | 0,0273 | 0,52 ^b | 2,74 ^a | 0,82 ^{ab} | 1,36 ^e | 171,37 |
| 473 | 0,0435 | 14,09 ^a | 16,21 ^a | 5,65 ^b | 12,01 ^d | 99,57 |
| 479 | 0,0115 | 54,76 ^a | 57,95 ^a | 32,63 ^b | 48,45 ^c | 50,08 |
| 485 | 0,0124 | 76,98 ^a | 77,68 ^a | 57,68 ^b | 70,78 ^b | 28,40 |
| 491 | 0,0467 | 96,03 ^a | 96,35 ^a | 89,98 ^b | 94,12 ^a | 7,92 |
| 497 | 0,0782 | 99,14 | 98,95 | 97,30 | 98,47 ^a | 2,33 |
| 503 | 0,9328 | 99,43 | 99,57 | 99,52 | 99,50 ^a | 0,93 |
| 515 | 1,0000 | 100,00 | 100,00 | 100,00 | 100,00 ^a | 0,00 |
| Breeder | -49,18 ^a | 49,98 ^a | 43,98 ^b | -94,12 | | |
| | P | | | | | |
| Breeder | 0,0001 | | | | | |
| Time | 0,0001 | | | | | |
| B x T | 0,0001 | | | | | |

Means followed by the same letter in the same row are not statistically different by the test of Tukey at 5% probability. Where: YB = young breeder (34 weeks); IB = intermediate-age breeder (44 weeks); OB = old breeder (72 weeks). Comparison of means within the same column.

Table 2 - Cumulative hatchings of chicks derived from broiler breeders of different ages (34, 44, and 72 weeks), % relative to total number of incubated eggs.

| Incubation (hours) | Breeder | | | | Incubation (mean) | CV (%) |
|--------------------|----------|---------------------|--------------------|--------------------|--------------------|--------|
| | P | YB | IB | OB | | |
| 449 | 1,0000 | 0,00 | 0,00 | 0,00 | 0,00 | 0,00 |
| 461 | 0,4946 | 0,00 | 0,19 | 0,17 | 0,12 ^e | 368,89 |
| 473 | 0,0213 | 12,24 ^{ab} | 14,68 ^a | 4,22 ^b | 10,38 ^d | 99,24 |
| 485 | 0,0001 | 68,79 ^a | 70,45 ^a | 42,44 ^b | 60,56 ^b | 31,84 |
| 497 | 0,0001 | 89,87 ^a | 89,07 ^a | 70,85 ^b | 83,26 ^a | 12,65 |
| 515 | 0,0001 | 89,86 ^a | 90,81 ^a | 72,70 ^b | 84,46 ^a | 11,65 |
| | P | | | | | |
| Breeder | 0,0001 | | | | | |
| B x T | 0,0001 | | | | | |

Means followed by the same letter in the same row are not statistically different by the test of Tukey at 5% probability. Where: YB = young breeder (34 weeks); IB = intermediate-age breeder (44 weeks); OB = old breeder (72 weeks). Comparison of means in the same column.

Vieira *et al.* (2005) also found that intermediate-age breeders' eggs started to hatch earlier than those laid by breeders with other ages, with the first hatchings occurring at 449 incubation hours - 12 hours earlier than those reported in the present study. Those authors

worked with multiple-stage incubators, and the heat generated by older embryos may have contributed to accelerate the development of a small percentage of eggs, thereby resulting in those very early hatchings. They also found the hatching distribution of old breeders' eggs was concentrated in the final periods of incubation, which is consistent with the findings of the present study. However, those authors observed late hatchings also for young breeders' eggs. Wilson (1991) and Marques (1994) also found a longer total incubation period for eggs from breeders at the onset of lay.

Hatching distribution, according to Wilson (1991), depends on breeder age, hatchable egg storage time and temperature, incubation temperature, egg weight, season, and chicken type. According to Campos (2000), the type of incubator - single or multiple stages - also influences hatching distribution. Multiple-stage equipment incubates embryos in different developmental stages, making embryos in later developmental stages interfere with the metabolism of embryos in earlier developmental stages. Salazar (2000) observed that high-capacity incubators also tend to present wider variation in total incubation time.

Meijerhof (2003) identified incubator air flow as a factor that may influence hatching distribution or total time required for all eggs to hatch. According to that author, incubators must have efficient air circulation in the internal environment in order to efficiently heat eggs during early embryo development and to remove the excessive heat produced by the embryos after day 10 of incubation. The efficiency of this process is reduced as incubator capacity increases.

Rosa & Ávila (2000) added incubation humidity as a factor that may contribute to variations in incubation time. According to those authors, eggs of breeders older than 50 weeks of age present high water content in the albumen, thin eggshell, and lower area to volume ratio, thereby hindering gas exchange. Therefore higher water loss (14 - 15%) than the recommended values (10 - 12%) is required to supply higher oxygen levels to the embryo, resulting in faster embryo growth and shorter incubation periods.

O'Connor (1984) correlated egg weight with incubation time: when egg weight doubles, incubation time increases 16%. This finding is consistent with the results of Vieira *et al.* (2005), who observed that the time required for hatching increased in heavy eggs as compared to lighter eggs of intermediate-age breeders. On the other hand, Pedroso *et al.* (2005) did not find significant correlations between egg weight and hatching time.



Baião *et al.* (2005) found that laying hour influences incubation time of broiler chicks. Those authors observed that chicks derived from eggs of the fifth daily egg collection hatched earlier than those from the first collection, which could be explained by the fact that the former remain longer in the hen's reproductive tract, and thereby submitted to higher temperature, accelerating embryo development after lay. These findings are not consistent with those of Fasenko *et al.* (1992), who observed that eggs from the first collection were heavier, indicating that their embryos were more developed.

All these variation factors contribute to the differences in the hatching distribution results reported in literature. In the present experiment, 71% of the eggs had hatched until 485 incubation hours, and 94% until 491 incubation hours. Reis *et al.* (1997) observed that 56% of the eggs had hatched at 485 hours and 94% at 490 hours, whereas Hudson *et al.* (2004) reported 60% hatching at 480 hours and 86% at 491 incubation hours.

In general, longer (Olsen, 1942; Williams *et al.*, 1951; Hudson *et al.*, 2004; Vieira *et al.*, 2005; Pedroso *et al.*, 2005), shorter (McNally & Byerly, 1936; Smith & Bohren, 1975; Shanawany, (1984); Burton & Tullet, 1985) or similar (Reis *et al.*, 1997) incubation periods as breeders aged were found in literature.

Long fasting periods stimulate the production of corticosterone, a hormone known for its strong inhibitor effect on immune cells. Early chick feeding stimulated the development of the bursa and the intensive production of lymphocytes (Bigot *et al.*, 2001).

Table 3 shows the relations between egg components and breeder age. Egg weight linearly increased with breeder age, whereas the albumen to egg weight ratio was not influenced by breeder age. Significant relations were recorded for yolk weight and eggshell weight, with higher relative yolk weight and lower relative eggshell weight for old breeders' eggs, while eggs from young and intermediate-age breeders did not present any differences in these parameters.

The higher eggshell percentage relative to egg weight, followed by higher albumen percentage in young breeders' eggs, and the higher yolk percentage in old breeders' eggs were also found by Tullet & Burton (1982), Vieira *et al.* (2005), Gomes *et al.* (2005a), and Ferreira *et al.* (2005).

Table 4 lists the relative data of the incubation waste analysis. The hatchability and the fertility of old breeders' eggs were lower than those of young and

intermediate-age breeders, which were not significantly different.

Table 3 - Egg weight and albumen, yolk, and eggshell percentage as a function of breeder age.

| Breeder | Egg weight, g | Albumen, % | Yolk, % | Eggshell, % |
|--------------|--------------------|------------|--------------------|--------------------|
| YB | 58.04 ^c | 57.20 | 28.75 ^b | 12.83 ^a |
| IB | 64.17 ^b | 57.33 | 29.84 ^b | 12.35 ^a |
| OB | 72.89 ^a | 56.58 | 31.59 ^a | 11.33 ^b |
| CV, % | 11.88 | 4.65 | 8.17 | 10.69 |
| P | 0.0001 | 0.6203 | 0.0005 | 0.0004 |

Genetic line: Isa-Vedette. Means followed by the same letter in the same column are not statistically different by the test of Tukey at 5% probability. Where: YB = young breeder (34 weeks); IB = intermediate-age breeder (44 weeks); OB = old breeder (72 weeks).

Table 4 - Embryo diagnosis of non hatched eggs laid by breeders with different ages, %.

| Breeder | Hatch Interm. | Hatchab. Late | Fert. PD | Inf. PL |
|--------------------|--------------------|--------------------|--------------------|--------------------|
| Young | 87,86 ^a | 93,46 ^a | 95,60 ^a | 4,40 ^b |
| 3,69 ^{ab} | 0,57 | 0,71 ^b | 0,14 | 0,43 |
| Interm. | 88,79 ^a | 93,63 ^a | 96,43 ^a | 3,57 ^b |
| 3,41 ^b | 0,32 | 1,46 ^{ab} | 0,32 | 0,32 |
| Old | 70,34 ^b | 86,30 ^b | 87,95 ^b | 12,04 ^a |
| 5,91 ^a | 0,68 | 3,18 ^a | 0,23 | 0,45 |
| CV, % | 12,22 | 4,29 | 4,46 | 69,55 |
| 48,09 | 113,43 | 107,21 | 205,20 | 167,76 |
| P | 0,0001 | 0,0001 | 0,0001 | 0,0001 |
| 0,0431 | 0,5651 | 0,0280 | 0,7710 | 0,9413 |

Where: Hatchb = hatchability, Fert. = fertility; Inf. = infertility; Early = early embryo mortality (1-7 days); Interm. = intermediate embryo mortality (8-14 days); Late = late embryo mortality (15-21 days); PD = pipped and dead; PL = pipped and alive. Means followed by the same letter in the same column are not statistically different by the test of Tukey at 5% probability.

Eggs from young and old breeders presented higher early embryo mortality percentages. Late embryo mortality was higher in old breeders' eggs. Eggs from intermediate-age breeders had the lowest general mortality rate. Live and dead pipped chick percentage and intermediate embryo mortality were not influenced by breeder age. In general, old breeder's eggs presented higher infertility and total embryo mortality (22.49%) as compared to young (9.94%) and intermediate-age breeders' eggs (9.40%). These findings are consistent with those reported by Mcdaniel *et al.* (1979), Wilson (1991), Reis *et al.* (1997), Lapão *et al.* (1999), and Hudson *et al.* (2004).

Understanding the physiological peculiarities of each breeder age, as well as embryo development will aid the design of new incubation management practices that can contribute to reduce total embryo mortality.

Table 5 presents egg weight loss during incubation and the relation between egg weight and chick weight



at hatching. No differences were found for the relation between egg weight loss during incubation and breeder age, with an average rate of 11%. This is not consistent with the results of Gomes *et al.* (2005b) and Michalsky *et al.* (2005ab), who observed higher humidity loss in old breeders' eggs.

Table 5 - Egg weight, humidity loss during incubation, chick weight at hatching and egg weight relative to chick weight at hatching as a function of breeder age.

| Breeder | EW, g | HL, % | WH, g | CRW, % |
|---------|--------------------|--------|--------------------|--------|
| Young | 58,00 ^c | 10,92 | 41,80 ^b | 72,09 |
| Interm. | 61,72 ^b | 10,80 | 43,70 ^b | 70,81 |
| Old | 66,28 ^a | 11,15 | 47,25 ^a | 72,45 |
| CV, % | 6,02 | 3,83 | 5,70 | 3,21 |
| P | 0,0017 | 0,7887 | 0,0233 | 0,7023 |

Where: PO = Egg weight; HL = humidity loss during incubation; WH = chick weight at hatching; CRW = egg weight relative to chick weight at hatching. Means followed by the same letter in the same column are not statistically different by the test of Tukey at 5% probability.

Chick weight at hatching was higher for those derived from old breeders' eggs, whereas no difference was found for those derived from young and intermediate-age breeders' eggs. Chick weight to egg weight ratio was not affected by breeder age, with an average of 72%. These data are consistent with those of Shanawany (1984) and Yannakopoulos (1988), but are different from the findings of Gomes *et al.* (2005a), who observed a higher ratio for chicks from young breeders. Campos *et al.* (2003) verified a 73% ratio between egg weight and chick weight, as did Vieira *et al.* (2005).

Table 6 presents the relations among chick weight, yolk-sac weight, and breeder age of chicks that hatched at 485 hours and that were removed from the hatcher at 515 hours. It is observed that 30 hours after hatching, chick weight continued to have a linear relation with breeder age. However, weight loss due to dehydration and to the utilization of yolk sac reserves changed: chicks derived from old and intermediate-age breeders lost more weight (average 5.56%) than those derived from young breeders (2.92%). As to yolk-sac weight at 485 hours, chicks derived from intermediate-age breeders used this reserve more (41.10%) than those derived from young and old breeders, which average utilization was 23%. Wyatt *et al.* (1985) and Nir & Levanon (1993) reported 5 to 32% body weight loss in chicks kept for 14 to 32 hours in the hatcher after hatching.

Breeder age influenced yolk sac weight relative to chick weight. Chicks from old breeders presented

higher relative yolk sac weight (16%) as compared to those from young breeders (10.41%). Chicks from intermediate-age breeders presented an intermediate value (13.43%), which was not different from young or old breeders. Vieira *et al.* (2005) observed that chicks from old breeders, and heavy chicks from intermediate-age breeders presented higher yolk sac weight (16.58%) as compared to chicks from young breeders or those derived from light eggs laid by intermediate-age breeders (14.88%). These data indicate that yolk-sac weight may be also be related to egg weight, and not only to breeder's age. These results are consistent with those of Gomes *et al.* (2005), who observed heavier yolk sacs in chicks derived from old breeders (16.19%) as compared to those derived from young (13.67%) and intermediate-age breeders (15.74%). Murakami *et al.* (1991) recorded a value of 10% for this relation, whereas Sklan & Noy (2000) observed 20%.

Table 6 - Live weight and relative yolk sac weight at hatching as a function of breeder age.

| Breeder | WA, g | WB, g | Diff., % | YSA, % | YSB, % | Diff., % |
|---------|---------------------|---------------------|----------|---------------------|--------|----------|
| Young | 39,73 ^c | 38,57 ^c | 2,92 | 10,41 ^b | 8,34 | 19,88 |
| Interm. | 44,87 ^{bA} | 42,23 ^{bB} | 5,88 | 13,43 ^{ab} | 7,91 | 41,10 |
| Old | 47,67 ^{aA} | 45,17 ^{aB} | 5,24 | 15,78 ^a | 11,50 | 27,12 |

WA = weight of chicks hatched at 485 hours; WB = weight of chicks hatched 485 hours and removed from the hatcher at 515 hours (30 hours after hatching); YSA = relative yolk sac weight at 485 hours; YSB = relative yolk sac weight of chicks hatched 485 hours and removed from the hatcher at 515 hours (30 hours after hatching). Means followed by the same letter in the same column are not statistically different by the test of Tukey at 5% probability.

CONCLUSIONS

In the present experiment, 71% of the chicks hatched at 485 incubation hours or less, and 94% had hatched at 491 hours. It is suggested that chicks should be removed at least once before 510 incubation hours, which is the commonly used incubation period, thereby reducing the period of time most birds are submitted to water and feed fasting.

Breeder age did not influence total time required for all eggs to hatch, but affected hatching distribution within this period.

Eggs laid by old breeders presented higher percentages of infertile eggs and total embryo mortality, resulting in a lower hatchability percentage for those birds.



REFERENCES

- Akaike H. Factor-analysis and AIC. *Psychometrika* 1987; 52(3):317-332.
- Baião LEC, Costa ES, Silva PL. Efeito do horário de postura sobre o período de nascimento de pintos de corte. *Revista Brasileira de Ciência Avícola* 2005; 3(supl):6.
- Bigot K, Tesseraud S, Taouis M. The relation of egg weight to chick weight at hatching. *Productions Animales* 2001; 14:219-230.
- Burton FG, Tullet SG. The effects of egg weight and shell porosity on the growth and water balance on the chicken embryo. *Comparative Biochemistry and Physiology* 1985; 81A:377-385.
- Campos EJ. *Avicultura: razões fatos e divergências*. Belo Horizonte: FEP-MVZ; 2000.
- Campos EJ, Santos JEC. O efeito de linhagens sobre o desenvolvimento embrionário. In: Macari M, Gonzales E, editores. *Manejo da incubação*. 2.ed. Jaboticabal: FACTA; 2003; p. 353-361.
- Dibner JJ, Knight CD, Kitchell ML, Atwell CA, Downs AC, Ivey FJ. Early feeding and the development of the immune system. *Journal of Applied Poultry Research* 1998; 7:425-436.
- Fanguy RC, Misra LK, Vo KV. Effect of delayed placement on growth performance of commercial broilers. *Poultry Science* 1980; 59:1215-1220.
- Fasenko GM, Hardin RT, Robinson FE. Variability in preincubation embryonic development in domestic fowl. *Poultry Science* 1992; 71:1374-1383.
- Gomes FS, Santos GCF, Silva PL. Efeito da linhagem e idade de reprodutoras pesadas na qualidade dos pintos de um dia. *Revista Brasileira de Ciência Avícola* 2005a; 3(supl):19.
- Gomes FS, Santos GCF, Silva PL. Efeito da linhagem e idade de reprodutoras pesadas na qualidade dos ovos incubáveis. *Revista Brasileira de Ciência Avícola* 2005b; 3(supl):20.
- Gonzales E, Kondo N, Saldanha ESPB, Loddy MM, Careghi C, Decuyper E. Performance and physiological parameters of broilers chickens subjected to fasting on the neonatal period. *Poultry Science* 2003; 82:1250-1256.
- Hager JE, Beane WL. Posthatch incubation time on early growth of broiler chicks. *Poultry Science* 1983; 62:247-254.
- Halevy O, Geyra A, Barak M, Uni Z, Sklan D. Early posthatch starvation decreases satellite cell proliferation and skeletal muscle growth in chicks. *Journal of Nutrition* 2000; 130:858-864.
- Hudson BP, Fairchild BD, Wilson JL, Dozier WA, Buhr RJ. Breeder age and zinc source in broiler breeder hen diets on progeny characteristics at hatching. *Journal of Applied Poultry Research* 2004; 13:55-64.
- Kingston DJ. Some hatchery factors involved in early chick mortality. *Australian Veterinary Journal* 1979; 55:418-421.
- Lapão C, Gama LT, Soares MC. Effects of broiler breeder age and length of egg storage on albumen characteristics and hatchability. *Poultry Science* 1999; 78:640-645.
- Marques, D. *Fundamentos básicos de incubação industrial*. 2.ed. São Paulo: CASP S/A; 1994.
- McDaniel GR, Roland DA, Coleman MA. The effect of eggs shell quality on hatchability and embryonic mortality. *Poultry Science* 1979; 58:10-13.
- McNally EH, Byerly TC. Variation in the development of embryos of hen eggs. *Poultry Science* 1936; 15:280-283.
- Meijerhof R. *Temperatura do embrião: uma nova ferramenta*. Ave Word 2003; 2.
- Michalsky VB, Cançado SV, Lara LJC, Baião NC, Santos GC, Lana AMQ. Efeito da umidade na incubação e da idade da matriz leve sobre as relações entre o coração, o saco vitelino e o peso do pinto. *Revista Brasileira de Ciência Avícola* 2005a; 3(supl):12.
- Michalsky VB, Cançado SV, Lara LJC, Baião NC, Santos GC, Lana AMQ. Efeito da umidade na incubação e da idade da matriz leve sobre a eclosão e parâmetros de ovos e pintos. *Revista Brasileira de Ciência Avícola* 2005b; 3 (supl):13.
- Murakami H, Akiba Y, Horiguchi M. Growth and utilization of nutrients in newly-hatched chicks with or without removal of residual yolk. *Grow, Development and Aging* 1991; 56:75-84.
- Nir I, Levanon M. Effect of posthatch holding time performance and residual yolk and liver composition. *Poultry Science* 1993; 72:1994-1997.
- Nitsan Z. The development of digestive tract in posthatched chicks. 10th European Symposium on Poultry Nutrition; 1995; Antalya. Turquia. p.21-28.
- Olsen MW. The effect of age and weight of turkey eggs on the length of incubation period. *Poultry Science* 1942; 21:532-535.
- O'Connor RJ. *The growth and development of birds*. New York: John Wiley & Sons; 1984.
- Pedroso AA, Stringhini JH, Leandro NSM, Xavier AS, Lima FG, Barbosa CE. Desempenho e biometria de órgãos digestórios de frangos provenientes de matrizes jovens após diferentes intervalos de alojamento. *Revista Brasileira de Ciência Avícola* 2005; 3(supl):5.
- Reis LH, Gama LT, Soares MC. Effects of short storage conditions and broiler age on hatchability, hatching time, and chick weights. *Poultry Science* 1997; 76:1459-1466.
- Rosa PS, Ávila VS. Variáveis relacionadas ao rendimento da incubação de ovos em matrizes de frangos de corte [comunicado técnico 246]. Concórdia (SC): Embrapa Suínos e Aves; 2000.
- Salazar AI. El proceso de incubación. *Avicultura Profesional* 2000;



18:26-30.

SAS Institute. SAS user's guide: Sstatistics. Cary, NC; 1998.

Schmidt GS, Figueiredo EAP, Ávila VS. Incubação: características dos ovos incubados. *Avicultura Industrial* 2003; 8:18-24.

Shanawany MM. Inter-relationship between egg weight, parental age and embryonic development. *British Poultry Science* 1984; 25: 449-455.

Sklan D, Noy Y, Hoyzman A, Rozemboim I. Decreasing weight loss in the hatchery by feeding chicks and poults in hatching trays. *Journal of Applied Poultry Research* 2000; 9:42-148.

Smith KP, Bohren BB. Age of pullets effect on hatching time, egg weight and hatchability. *Poultry Science* 1975; 54:959-963.

Vieira SL, Almeida JG, Lima AR, Conde ORA, Olmos AR. Hatching distribution of eggs varying in weight and breeder age. *Brazilian Journal of Poultry Science* 2005; 7:73-78.

Vieira SL, Pophal S. Nutrição pós-eclosão em frangos de corte. *Revista Brasileira de Ciência Avícola* 2000; 2:189-286.

Williams C, Godfrey GF, Thompson RB. The effect of rapidity of hatching on growth, egg production, mortality and sex ratio in the domestic fowl. *Poultry Science* 1951; 30:599-606.

Wilson HR. Interrelationships of egg size, chick size, posthatching growth and hatchability. *World's Poultry Science Journal* 1991; 47:5-20.

Wyatt CL, Weaver WD, Beane WL. Influence of egg size, egg shell quality, and posthatch holding time on broiler performance. *Poultry Science* 1985; 64:2049-2055.

Yannakopoulos AL. Lack of effect of ovoposition time and parental age on chick weight when egg weight remains constant. *British Poultry Science* 1988; 29:431-434.