

Research Article

Riluzole Stimulates BDNF Release from Human Platelets

Patrick Türck and Marcos Emilio Frizzo

Department of Morphological Sciences, Institute of Basic Health Sciences, Federal University of Rio Grande do Sul, Rua Sarmento Leite, 500, 90050-170 Porto Alegre, RS, Brazil

Correspondence should be addressed to Marcos Emilio Frizzo; frizzo@ufrgs.br

Received 1 July 2014; Accepted 17 September 2014

Academic Editor: Gianluca Coppola

Copyright © 2015 P. Türck and M. E. Frizzo. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Brain-derived neurotrophic factor (BDNF) has several functions in the central nervous system, where it contributes to brain development and its functionality through affecting neuronal survival and activity and also modulating neurotransmitter levels. This neurotrophin is also found in the serum, but its origin and peripheral function remain unknown. Although the source of circulating BDNF is uncertain, it is stored in platelets and can be released through pharmacological treatment. Decreased levels of BDNF in the serum have been related to the pathophysiology of depression, and this relationship is reinforced by the reversal of this condition by treatment with antidepressants. Recently, riluzole has been proposed for the treatment of depression because it has the ability to lower extracellular glutamate levels and increase BDNF expression; and both mechanisms could be associated with its antidepressant action. Considering that riluzole enhances BDNF levels in the serum of patients, we investigated if treatment with this drug could stimulate the release of this neurotrophin from human platelets obtained from healthy subjects. When platelets were incubated with riluzole for 4 h, the basal value of BDNF ($92.9 \pm 11.1 \text{ pg } 10^{-6}$ platelets) was significantly increased ($P < 0.05$, $n = 27$). This stimulatory effect was achieved at low concentrations of riluzole (from $10 \mu\text{M}$) and was not observed when platelets were incubated with the drug for 24 h. The direct action of riluzole evoking BDNF release from human platelets at therapeutic concentrations is important and may contribute to the understanding of its mechanisms of action in the treatment of depression.

1. Introduction

Brain-derived neurotrophic factor (BDNF) contributes to brain development [1, 2] and is related to neuronal survival and activity since it acts as a modulator of neurotransmitter levels and participates in neuronal plasticity [3, 4]. In the human, monkey, and rat, BDNF is also found in the serum at significant levels [5–7], but the origin and function of this neurotrophin remain unknown. Investigators have mentioned the brain as the source for this circulating neurotrophin [8], even though it has been demonstrated that BDNF crosses the blood-brain barrier (BBB) in both directions [9, 10]. Indeed, BDNF may originate from neurons and glia cells [9, 10]; however, it is also released at significant rates by other peripheral tissues, such as different epithelia, where its amounts may reach levels higher than those found in the central nervous system (CNS) [11]. Other examples of BDNF origins other than from CNS are white cells [12–14] and platelets; the latter contain significant quantities of

this protein and might provide an important source of this circulating neurotrophin [5]. It has been shown that more than 99% of blood BDNF proteins are stored in platelets and that these proteins can be released into the serum [6] through pharmacological treatment [15, 16].

Recent studies have reported changes in serum BDNF levels in patients with psychiatric diseases [17–20], such as major depressive disorders [21]. The relationship between decreased BDNF levels and the pathophysiology of depression is supported by several reports [21–25]. Pandey et al. [26] showed that gene expression of BDNF in lymphocytes and its protein expression in platelets from adult and pediatric depressed patients were significantly decreased, and the authors proposed that it could be a target for antidepressant drugs. In fact, some antidepressants increase BDNF expression [27] and also may evoke BDNF release from platelets, in a dose-dependent manner after direct treatment *in vitro* [15]. The BDNF concentration in the serum increases after intravenous treatment with an antidepressant, and the effect

of these drugs on BDNF release from platelets was related to the level of this neurotrophin in the peripheral blood [15].

Recently, glutamatergic modulators have been proposed as a strategy for the treatment of mood disorders [28]. Among the drugs proposed is riluzole (2-amino-6-trifluoromethoxy benzothiazole), which was originally developed as an anti-convulsant [29] but has been used in a number of trials for psychiatric conditions in which glutamate excess has been proposed as part of the pathologic mechanism [30–33]. Different mechanisms of action have been reported for riluzole [32], which probably explains its complex pharmacological effects. For instance, a stimulatory effect on glutamate uptake was observed at low glutamate concentration [34], and this ability to lower extracellular glutamate levels was suggested as its mechanism of antidepressant action, at least partially [32]. However, other mechanisms cannot be ruled out, since riluzole also increases the BDNF expression [35, 36], which could also contribute to its antidepressant action [32]. Treatment with riluzole significantly increases serum levels of BDNF in patients [37]. Considering that BDNF in the blood is thought to originate from platelets and is evoked by antidepressant drugs, we decided to investigate if riluzole could stimulate the release of this neurotrophin from human platelets.

2. Methods

2.1. Subjects. Human blood was collected from 27 healthy male volunteers registered as donors in the Hemotherapy Service of the Clinical Hospital of Porto Alegre, Rio Grande do Sul, Brazil. The study was carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans. Informed consent was obtained from the donors and their privacy rights were observed.

2.2. Procedures. Two samples of 4 mL were taken from the antecubital vein of each donor and placed in vacutainers (BD Franklin Lakes, USA) containing K_3 -EDTA. Immediately, they were gently inverted 10 times and placed in an ABX Micros ES 60 hematology analyzer (HORIBA ABX SAS, Japan) to determine the number of platelets in each blood sample. Platelets were isolated as previously described by Mangano and Schwarcz [38]. The vacutainers were then centrifuged at $300 \times g$ for 5 min at $4^\circ C$ and the platelet-rich plasma (PRP) was obtained. The supernatant (PRP) was carefully removed with a plastic pipette tip, with care not to disturb the leukocyte layer. The volume of PRP collected from each sample was recorded and the PRP was transferred to a microcentrifuge tube. The PRP was then centrifuged at $7000 \times g$ for 10 min at $4^\circ C$. The plasma was discarded and the pellet was resuspended in 0.5 mL of 0.32 M phosphate-buffered sucrose (pH 7.4 at $4^\circ C$). The suspension, hereafter referred to as the platelet concentrate (PC), was repeatedly passed through a plastic pipette tip of 1 mL until the visible platelet aggregates were eliminated. An additional 0.5 mL of buffered sucrose was added to the suspension and the solution was mixed with 5 gentle inversions. The PC was again centrifuged at $7000 \times g$ for 5 min at $4^\circ C$. The supernatant was

discarded, and the pellet was resuspended in a volume of 0.32 M phosphate-buffered sucrose (pH 7.4 at $4^\circ C$), equal to one-fifth of the initial volume of PRP obtained. After that, the two PC suspensions obtained from each subject were blended and the number of platelets was determined again. As previously observed by our group, the platelets should be counted in the PC, after the platelets are washed [39]. The mean platelet volume (MPV) and platelet distribution width (PDW) were also determined in both the whole blood and the PC to evaluate potential variations in the platelets during the processing. The incubation was performed in a 96-well plate; to each well were added 20×10^6 platelets in 130 μL of Tris-citrate buffer (112.8 mM NaCl, 4.5 mM KCl, 1.1 mM KH_2PO_4 , 1.1 mM $MgSO_4$, 11 mM Na_3 -citrate, 25 mM Tris-HCl, and 10.2 mM glucose), pH 6.5. Drugs were diluted in Tris-citrate buffer, and the final volume in the well was 150 μL .

2.3. BDNF Protein Assay. BDNF levels in the supernatants were measured using a ChemiKine Brain Derived Neurotrophic Factor Sandwich ELISA kit (Millipore, USA) following the manufacturer's instructions. All BDNF measurements were performed in triplicate on 96-well plates, and a standard curve was calculated for each experiment. Samples from the supernatants were diluted 1:16 in phosphate buffer solution (pH 7.4) for BDNF measurement. The platelet BDNF content was calculated by dividing the result for BDNF obtained by the total platelet count from the same individual and was expressed as pg BDNF 10^{-6} platelets. The optical density of each well was measured using a microplate reader (EZ Read 400, Biochrom, UK) set to 450 nm; the optical densitometry data were analyzed with the software Galapagos (Biochrom, UK). The sensitivity was 7.8 pg BDNF mL^{-1} and the assay exhibited no cross-reactivity with other members of the nerve growth factor family.

2.4. MTT Assay. The MTT assay was utilized to determine the effect of riluzole or sertraline on platelet viability after 4 or 24 h of drug exposure. For shorter incubation times (4 h), 20 μL MTT (5 mg mL^{-1}) was added to each well at the time that the platelets were plated, and it was maintained at room temperature by 4 h. For longer incubation times (24 h), 20 μL MTT (5 mg mL^{-1}) was added to each well 20 h after plating and maintained at room temperature for an additional 4 h. Subsequently, 150 μL of DMSO was added to dissolve the formazan, which was detected using a microplate reader (EZ Read 400, Biochrom, UK). The absorbance was read at 570 nm; the value obtained (UAbs) was expressed as UAbs $20 \cdot 10^{-6}$ platelets.

2.5. Statistical Methods. Values are reported as mean \pm SEM, and the statistical analysis was conducted using SPSS. The data were normally distributed, as determined by the Shapiro-Wilk normality test, and were analyzed through one-way ANOVA (for BDNF) or two-way ANOVA (for MTT) followed by Tukey's multiple comparisons test (alpha at 0.05).




FIGURE 1: Scatter plot of BDNF released from human platelets. The mean of BDNF values (92.9 ± 11.1 pg 10^{-6} platelets) is indicated by the dotted bar ($n = 27$).




FIGURE 2: Stimulatory effect of riluzole on BDNF release. Platelets obtained from healthy donors were incubated with different concentrations of riluzole (1, 10, 40, or 100 μ M) for 4 h. Control is represented by the dotted line. Data are presented as mean \pm SEM. * Different from control ($n = 27$, $P < 0.05$).

3. Results

We used washed platelets to test if riluzole could stimulate BDNF release, acting directly on these cells. Initially, the platelet indices (cell count, MPV, and PDW) were measured in whole blood and again after obtaining the PC, to evaluate if these parameters changed during the processing. Then, the platelet number quantified in each PC was used to normalize the BDNF quantity in the respective experiment. Our data showed that the platelet count should be determined after the PC is obtained, since the cell number was reduced during the process (yield of $42 \pm 8\%$). Analysis of PC showed an absence of contaminant cells; and despite the loss of platelets, the MPV and PDW indices were not changed after the cells were obtained. The platelet indices MPV and PDW in the whole blood were $8.1 \pm 0.6 \mu\text{m}^3$ and $14.9 \pm 1.0\%$, respectively. When compared with PC, we did not observe significant differences for MPV and PDW, which showed levels of $8.2 \pm 0.6 \mu\text{m}^3$ and $15.5 \pm 1.2\%$, respectively. In our analysis, the platelet number achieved in the PC was $1545 \pm 338 \times 10^3 \text{ mm}^{-3}$.

In our experimental conditions, the basal values of BDNF released from platelets of donors ranged from 9.0 to

220.2 pg 10^{-6} platelets. The wide distribution of BDNF quantified in the study group is depicted in Figure 1.

Given the recent evidence that riluzole treatment causes a significant increase of BDNF in the serum of patients [37], the effect of this drug on BDNF release from human platelets was tested. Platelets from healthy volunteers were treated with different concentrations of riluzole for 4 or 24 h at room temperature. When platelets were incubated with riluzole for 4 h, the basal value of BDNF as quantified from the controls (92.9 ± 11.1 pg 10^{-6} platelets) was significantly increased ($P < 0.05$). Even for platelets from donors who showed lower basal levels of BDNF, treatment with riluzole stimulated the release of this neurotrophin. The increase mediated by riluzole was significant beginning with 10 μ M (15%) and was maintained up to 40 μ M (22%) and 100 μ M (20%). We also tested the effect of 1 μ M riluzole, which did not differ from the control (Figure 2).

The effect of riluzole on the BDNF release determined in our model was reproducible in repeated runs, even though it varied from individual to individual. Considering this variability in the basal levels of BDNF among different donors (Figure 1), in each set of experiments, the drug effect was




FIGURE 3: Platelet viability during the incubation period. The effect of different concentrations of riluzole on platelet viability was determined at 4 or 24 h of incubation. No significant differences were observed among the groups. Data are presented as mean \pm SEM ($n = 27$).

compared with the respective control. We also used sertraline ($0.3 \mu\text{M}$) as a positive control, since it was recently reported to be a potent inducer of BDNF release from platelets [15]. However, contrary to expectations, in our experiments, the sertraline did not significantly stimulate BDNF release (data not shown).

The increase in the release of BDNF evoked by riluzole was not observed when platelets were incubated with the drug for 24 h. In order to determine if platelet viability was affected by riluzole during the exposure, we used the MTT assay. Hence, the activity of NAD(P)H-dependent cellular oxidoreductase enzymes was evaluated after 4 and 24 h incubation. The viability of the untreated platelets did not differ after the two incubation periods, although slightly less formazan was produced after 24 h (Figure 3).

4. Discussion

We evaluated the platelet parameters MPV and PDW, since these indices have been reported to correlate with platelet function [40]. MPV is a measurement that is commonly used to describe platelet size and is an indicator of activated platelets [41, 42], while PDW represents the range of variability in platelet volume [43]. Taking into account that MPV and PDW did not change before the platelets were obtained and after they were processed, we can state that the platelets used in this study were not activated at the time of exposure to riluzole. Importantly, we did not select a subpopulation of these cells, since no significant difference was observed in the PDW. Another important point is that our protocol used only platelets, and, consequently, the BDNF quantified cannot be attributed to contaminant cells such as leukocytes.

In our study, we observed a direct action of riluzole, evoking BDNF release from human platelets. The stimulatory effect was achieved at low concentrations (from $10 \mu\text{M}$), which could be important information for the clinical use

of riluzole. Although this increase was not as large as found for another antidepressant [15], the stimulatory effect was reproducible when compared with the respective controls, despite the variations among different donors. The absence of a sertraline effect on BDNF release might be due to our use of human cells, whereas Watanabe et al. [15] used rat platelets. Unfortunately, Watanabe and coauthors did not report the basal values of BDNF that they obtained, which would allow comparison with our data.

Riluzole evoked an acute effect on the platelets, which was not observed after longer incubation times. The release of BDNF in response to riluzole acutely (4 h) and not later (24 h) suggests that its effect derived from evoking neurotrophin release from the platelet pool and is not related to a stimulatory effect on neurotrophin synthesis. This is in accordance with data showing that mRNA expression of BDNF in human platelets is extremely low [5, 10].

The novel finding that riluzole elicits BDNF release from human platelets is important, since this situation may occur peripherally and also in deep regions of the CNS, where platelets and astrocytes are very close and where this neurotrophin is able to pass through the BBB. Beyond the peripheral consequences of this release mediated by riluzole, the effects of this neurotrophin on the CNS may be complex and significant, especially regarding the glutamatergic system. It has been shown that BDNF exerts acute effects on glutamatergic synaptic transmission and plasticity, that is, enhancing excitatory synaptic transmission through pre- and postsynaptic mechanisms [44]. On the other hand, its stimulatory effect on the expression of astroglial glutamate transporters and the consequent increase in glutamate uptake capacity has also been described [45]. More recently, it was demonstrated that BDNF upregulates the protein expression of the vesicular glutamate transporters (VGLUT1 and VGLUT2) in hippocampal neurons [46], which reinforces the participation of BDNF as a modulator of the glutamatergic synapse.

Clinically, riluzole has been used in trials for psychiatric conditions where glutamate excess is proposed as part of the pathologic mechanism [47]. It is also suggested that it produces antidepressant and anxiolytic effects in the treatment of resistant depression [32]. Its effect in these conditions is associated with the ability to reduce extracellular glutamate levels and also may involve its stimulatory action on BDNF expression [32]. Therefore, the demonstration that riluzole causes BDNF release at low concentrations is significant *in vivo* and shows the importance of studying platelets from patients treated with this drug. Studies to clarify the mechanisms related to BDNF release in human platelets are currently in progress in our laboratory.

5. Conclusions

The new effect described for riluzole may contribute to the understanding of the mechanisms involved with its therapeutic action, reinforcing the suggestion for its use in psychiatry, such as in the treatment of depression.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publishing of this paper.







Acknowledgment

This research was supported by grants from the Brazilian National Research Council (CNPq).

References

- [1] G. R. Lewin and Y.-A. Barde, "Physiology of the neurotrophins," *Annual Review of Neuroscience*, vol. 19, pp. 289–317, 1996.
- [2] S. D. Croll, N. Y. Ip, R. M. Lindsay, and S. J. Wiegand, "Expression of BDNF and trkB as a function of age and cognitive performance," *Brain Research*, vol. 812, no. 1-2, pp. 200–208, 1998.
- [3] E. Castrén, M. Pitkänen, J. Sirviö et al., "The induction of LTP increases BDNF and NGF mRNA but decreases NT-3 mRNA in the dentate gyrus," *NeuroReport*, vol. 4, no. 7, pp. 895–898, 1993.
- [4] J. L. Jankowsky and P. H. Patterson, "Cytokine and growth factor involvement in long-term potentiation," *Molecular and Cellular Neurosciences*, vol. 14, no. 6, pp. 273–286, 1999.
- [5] H. Yamamoto and M. E. Gurney, "Human platelets contain brain-derived neurotrophic factor," *Journal of Neuroscience*, vol. 10, no. 11, pp. 3469–3478, 1990.
- [6] S. F. Radka, P. A. Holst, M. Fritsche, and C. A. Altar, "Presence of brain-derived neurotrophic factor in brain and human and rat but not mouse serum detected by a sensitive and specific immunoassay," *Brain Research*, vol. 709, no. 1, pp. 122–301, 1996.
- [7] T. Mori, K. Shimizu, and M. Hayashi, "Levels of serum brain-derived neurotrophic factor in primates," *Primates*, vol. 44, no. 2, pp. 167–169, 2003.
- [8] R. Katoh-Semba, R. Wakako, T. Komori et al., "Age-related changes in BDNF protein levels in human serum: differences between autism cases and normal controls," *International Journal of Developmental Neuroscience*, vol. 25, no. 6, pp. 367–372, 2007.
- [9] W. Pan, W. A. Banks, M. B. Fasold, J. Bluth, and A. J. Kastin, "Transport of brain-derived neurotrophic factor across the blood-brain barrier," *Neuropharmacology*, vol. 37, no. 12, pp. 1553–1561, 1998.
- [10] F. Karege, M. Schwald, and M. Cisse, "Postnatal developmental profile of brain-derived neurotrophic factor in rat brain and platelets," *Neuroscience Letters*, vol. 328, no. 3, pp. 261–264, 2002.
- [11] M. Lommatzsch, A. Braun, A. Mannsfeldt et al., "Abundant production of brain-derived neurotrophic factor by adult visceral epithelia: implications for paracrine and target-derived neurotrophic functions," *The American Journal of Pathology*, vol. 155, no. 4, pp. 1183–1193, 1999.
- [12] A. Braun, M. Lommatzsch, A. Mannsfeldt et al., "Cellular sources of enhanced brain-derived neurotrophic factor production in a mouse model of allergic inflammation," *The American Journal of Respiratory Cell and Molecular Biology*, vol. 21, no. 4, pp. 537–546, 1999.
- [13] A. Gielen, M. Khademi, S. Muhallab, T. Olsson, and F. Piehl, "Increased brain-derived neurotrophic factor expression in white blood cells of relapsing-remitting multiple sclerosis patients," *Scandinavian Journal of Immunology*, vol. 57, no. 5, pp. 493–497, 2003.
- [14] M. Kerschensteiner, E. Gallmeier, L. Behrens et al., "Activated human T cells, B cells, and monocytes produce brain-derived neurotrophic factor in vitro and in inflammatory brain lesions: a neuroprotective role of inflammation?" *Journal of Experimental Medicine*, vol. 189, no. 5, pp. 865–870, 1999.
- [15] K. Watanabe, E. Hashimoto, W. Ukai et al., "Effect of antidepressants on brain-derived neurotrophic factor (BDNF) release from platelets in the rats," *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, vol. 34, no. 8, pp. 1450–1454, 2010.
- [16] P. Stoll, A. Plessow, K. Bratke, J. C. Virchow, and M. Lommatzsch, "Differential effect of clopidogrel and aspirin on the release of BDNF from platelets," *Journal of Neuroimmunology*, vol. 238, no. 1-2, pp. 104–106, 2011.
- [17] K. Toyooka, K. Asama, Y. Watanabe et al., "Decreased levels of brain-derived neurotrophic factor in serum of chronic schizophrenic patients," *Psychiatry Research*, vol. 110, no. 3, pp. 249–257, 2002.
- [18] K. B. Nelson, J. K. Grether, J. M. Dambrosia et al., "Neuropeptides and neurotrophins in neonatal blood of children with autism or mental retardation," *Annals of Neurology*, vol. 49, no. 5, pp. 597–606, 2001.
- [19] K. Miyazaki, N. Narita, R. Sakuta et al., "Serum neurotrophin concentrations in autism and mental retardation: a pilot study," *Brain & Development*, vol. 26, no. 5, pp. 292–295, 2004.
- [20] K. Hashimoto, Y. Iwata, K. Nakamura et al., "Reduced serum levels of brain-derived neurotrophic factor in adult male patients with autism," *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, vol. 30, no. 8, pp. 1529–1531, 2006.
- [21] E. Shimizu, K. Hashimoto, N. Okamura et al., "Alterations of serum levels of brain-derived neurotrophic factor (BDNF) in depressed patients with or without antidepressants," *Biological Psychiatry*, vol. 54, no. 1, pp. 70–75, 2003.
- [22] F. Karege, G. Perret, G. Bondolfi, M. Schwald, G. Bertschy, and J.-M. Aubry, "Decreased serum brain-derived neurotrophic factor levels in major depressed patients," *Psychiatry Research*, vol. 109, no. 2, pp. 143–148, 2002.
- [23] F. Karege, G. Bondolfi, N. Gervasoni, M. Schwald, J.-M. Aubry, and G. Bertschy, "Low Brain-Derived Neurotrophic Factor

- (BDNF) levels in serum of depressed patients probably results from lowered platelet BDNF release unrelated to platelet reactivity," *Biological Psychiatry*, vol. 57, no. 9, pp. 1068–1072, 2005.
- [24] A. B. M. Cunha, B. N. Frey, A. C. Andreazza et al., "Serum brain-derived neurotrophic factor is decreased in bipolar disorder during depressive and manic episodes," *Neuroscience Letters*, vol. 398, no. 3, pp. 215–219, 2006.
- [25] R. Machado-Vieira, M. O. Dietrich, R. Leke et al., "Decreased plasma brain derived neurotrophic factor levels in unmedicated bipolar patients during manic episode," *Biological Psychiatry*, vol. 61, no. 2, pp. 142–144, 2007.
- [26] G. N. Pandey, Y. Dwivedi, H. S. Rizavi, X. Ren, H. Zhang, and M. N. Pavuluri, "Brain-derived neurotrophic factor gene and protein expression in pediatric and adult depressed subjects," *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, vol. 34, no. 4, pp. 645–651, 2010.
- [27] C. Pittenger and R. S. Duman, "Stress, depression, and neuroplasticity: a convergence of mechanisms," *Neuropsychopharmacology*, vol. 33, no. 1, pp. 88–109, 2008.
- [28] C. Zarate Jr., R. MacHado-Vieira, I. Henter, L. Ibrahim, N. Diazgranados, and G. Salvadore, "Glutamatergic modulators: the future of treating mood disorders?" *Harvard Review of Psychiatry*, vol. 18, no. 5, pp. 293–303, 2010.
- [29] J. Mizoule, B. Meldrum, M. Mazadier et al., "2-Amino-6-trifluoromethoxy benzothiazole, a possible antagonist of excitatory amino acid neurotransmission—I: anticonvulsant properties," *Neuropharmacology*, vol. 24, no. 8, pp. 767–773, 1985.
- [30] S. J. Mathew, K. Keegan, and L. Smith, "Glutamate modulators as novel interventions for mood disorders," *Revista Brasileira de Psiquiatria*, vol. 27, no. 3, pp. 243–248, 2005.
- [31] S. J. Mathew, J. W. Murrough, M. Aan Het Rot, K. A. Collins, D. L. Reich, and D. S. Charney, "Riluzole for relapse prevention following intravenous ketamine in treatment-resistant depression: A pilot randomized, placebo-controlled continuation trial," *International Journal of Neuropsychopharmacology*, vol. 13, no. 1, pp. 71–82, 2010.
- [32] G. Sanacora, S. F. Kendell, Y. Levin et al., "Preliminary evidence of riluzole efficacy in antidepressant-treated patients with residual depressive symptoms," *Biological Psychiatry*, vol. 61, no. 6, pp. 822–825, 2007.
- [33] C. A. Zarate Jr., J. L. Payne, J. Quiroz et al., "An open-label trial of riluzole in patients with treatment-resistant major depression," *The American Journal of Psychiatry*, vol. 161, no. 1, pp. 171–174, 2004.
- [34] M. E. Frizzo, L. P. Dall'Onder, K. B. Dalcin, and D. O. Souza, "Riluzole enhances glutamate uptake in rat astrocyte cultures," *Cellular and Molecular Neurobiology*, vol. 24, no. 1, pp. 123–128, 2004.
- [35] R. Katoh-Semba, T. Asano, H. Ueda et al., "Riluzole enhances expression of brain-derived neurotrophic factor with consequent proliferation of granule precursor cells in the rat hippocampus," *The FASEB Journal*, vol. 16, no. 10, pp. 1328–1330, 2002.
- [36] I. Mizuta, M. Ohta, K. Ohta, M. Nishimura, E. Mizuta, and S. Kuno, "Riluzole stimulates nerve growth factor, brain-derived neurotrophic factor and glial cell line-derived neurotrophic factor synthesis in cultured mouse astrocytes," *Neuroscience Letters*, vol. 310, no. 2-3, pp. 117–120, 2001.
- [37] F. Squitieri, S. Orobello, M. Cannella et al., "Riluzole protects Huntington disease patients from brain glucose hypometabolism and grey matter volume loss and increases production of neurotrophins," *European Journal of Nuclear Medicine and Molecular Imaging*, vol. 36, no. 7, pp. 1113–1120, 2009.
- [38] R. M. Mangano and R. Schwarcz, "The human platelet as a model for the glutamatergic neuron: platelet uptake of L-glutamate," *Journal of Neurochemistry*, vol. 36, no. 3, pp. 1067–1076, 1981.
- [39] D. S. Rocha, S. K. Souza, T. G. H. Onsten, R. S. M. Silva, and M. E. Frizzo, "A simple method to quantify glycogen from human platelets," *Journal of Cytology & Histology*, vol. 5, article 217, 2014.
- [40] Q. Niu, R. Zhang, M. Zhao et al., "Differences in platelet indices between healthy Han population and Tibetans in China," *PLoS ONE*, vol. 8, no. 6, Article ID e67203, 2013.
- [41] A. Arikanoğlu, Y. Yücel, A. Acar et al., "The relationship of the mean platelet volume and C-reactive protein levels with mortality in ischemic stroke patients," *European Review for Medical and Pharmacological Sciences*, vol. 17, no. 13, pp. 1774–1777, 2013.
- [42] G. De Luca, G. G. Secco, M. Verdoia et al., "Combination between mean platelet volume and platelet distribution width to predict the prevalence and extent of coronary artery disease: results from a large cohort study," *Blood Coagulation and Fibrinolysis*, vol. 25, no. 1, pp. 86–91, 2014.
- [43] R. Ozdemir, C. Karadeniz, O. Doksoz et al., "Are mean platelet volume and platelet distribution width useful parameters in children with acute rheumatic carditis?" *Pediatric Cardiology*, vol. 35, no. 1, pp. 53–56, 2014.
- [44] J.-L. Martin and C. Finsterwald, "Cooperation between BDNF and glutamate in the regulation of synaptic transmission and neuronal development," *Communicative & Integrative Biology*, vol. 4, no. 1, pp. 14–16, 2011.
- [45] A. Rodriguez-Kern, M. Gegelashvili, A. Schousboe, J. Zhang, L. Sung, and G. Gegelashvili, "Beta-amyloid and brain-derived neurotrophic factor, BDNF, up-regulate the expression of glutamate transporter GLT-1/EAAT2 via different signaling pathways utilizing transcription factor NF- κ B," *Neurochemistry International*, vol. 43, no. 4-5, pp. 363–370, 2003.
- [46] C. V. Melo, M. Mele, M. Curcio, D. Comprido, C. G. Silva, and C. B. Duarte, "BDNF regulates the expression and distribution of vesicular glutamate transporters in cultured hippocampal neurons," *PLoS ONE*, vol. 8, no. 1, Article ID e53793, 2013.
- [47] P. Grant, J. Y. Song, and S. E. Swedo, "Review of the use of the glutamate antagonist riluzole in psychiatric disorders and a description of recent use in childhood obsessive-compulsive disorder," *Journal of Child and Adolescent Psychopharmacology*, vol. 20, no. 4, pp. 309–315, 2010.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

