



Plant responses to stresses: Role of ascorbate peroxidase in the antioxidant protection

Andréia Caverzan¹, Gisele Passaia¹, Silvia Barcellos Rosa², Carolina Werner Ribeiro², Fernanda Lazzarotto¹ and Márcia Margis-Pinheiro^{1,2*}

¹Programa de Pós-Graduação em Genética e Biologia Molecular, Departamento de Genética, Universidade Federal Rio Grande do Sul, Porto Alegre, RS, Brazil.

²Programa de Pós-Graduação em Biologia Celular e Molecular, Centro de Biotecnologia, Universidade Federal Rio Grande do Sul, Porto Alegre, RS, Brazil.

Abstract

When plants are exposed to stressful environmental conditions, the production of Reactive Oxygen Species (ROS) increases and can cause significant damage to the cells. Antioxidant defenses, which can detoxify ROS, are present in plants. A major hydrogen peroxide detoxifying system in plant cells is the ascorbate-glutathione cycle, in which, ascorbate peroxidase (APX) enzymes play a key role catalyzing the conversion of H₂O₂ into H₂O, using ascorbate as a specific electron donor. Different APX isoforms are present in distinct subcellular compartments, such as chloroplasts, mitochondria, peroxisome, and cytosol. The expression of APX genes is regulated in response to biotic and abiotic stresses as well as during plant development. The APX responses are directly involved in the protection of plant cells against adverse environmental conditions. Furthermore, mutant plants APX genes showed alterations in growth, physiology and antioxidant metabolism revealing those enzymes involvement in the normal plant development.

Keywords: ascorbate peroxidase, antioxidant system, reactive oxygen species, abiotic stress, mutant plants.

Plant Responses to Stresses

The exposure of plants to unfavorable environmental conditions increases the production of reactive oxygen species (ROS) such as, singlet oxygen (¹O₂), superoxide (O₂^{•-}), hydrogen peroxide (H₂O₂), and hydroxyl radical (OH[•]). The ROS detoxification process in plants is essential for the protection of plant cells and their organelles against the toxic effect of these species (Apel and Hirt, 2004; Mittler, 2002). The differences in subcellular localization and biochemical properties of antioxidant enzymes and the distinct responses in gene expression, in addition to the presence of non-enzymatic mechanisms, result in a versatile and flexible antioxidant system able to control the optimum ROS levels (Vranova *et al.*, 2002). The ROS detoxification systems include enzymatic and non-enzymatic antioxidant components (Scandalios, 2005). Ascorbate (AsA) and glutathione (GSH), non-enzymatic antioxidants are crucial for plant defense against oxidative stress, playing a key role as antioxidant buffers (Foyer and Noctor, 2005; Mittler, 2002). Other non-enzymatic antioxidants involved include

flavonoids, phenolic compounds, alkaloids, tocopherol and carotenoids (Gratão *et al.*, 2005).

Enzymatic antioxidants comprise superoxide dismutase (SOD), ascorbate peroxidase (APX), catalase (CAT), glutathione peroxidase (GPX) and peroxiredoxin (PrxR). These enzymes are present in practically all subcellular compartments. Usually, an organelle has more than one enzyme able to scavenge a single ROS (Mittler, 2002; Mittler *et al.*, 2004; Scandalios, 2005). The main hydrogen peroxide-detoxification system in plant chloroplasts is the ascorbate-glutathione cycle, in which APX is a key enzyme (Asada, 1992). APX utilizes AsA as specific electron donor to reduce H₂O₂ to water. The importance of APX and ascorbate-glutathione cycle is not restricted to chloroplasts; it also plays a role in ROS scavenging in cytosol, mitochondria and peroxisomes (Asada, 1992, 1999; Mittler *et al.*, 2004; Noctor and Foyer, 1998; Shigeoka *et al.*, 2002). The ROS-scavenging enzymes in plants have been widely studied and the results have demonstrated that, in response to environmental stress, APX activity generally increases along with other enzymes activities, such as CAT, SOD, and GSH reductase (Shigeoka *et al.*, 2002). Over the past ten years substantial efforts have been made to understand plant antioxidant system mechanisms (Figure 1). The in-

Send correspondence to Marcia Margis-Pinheiro. Departamento de Genética, Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves 9500, 91501-970 Porto Alegre, RS, Brazil. E-mail: marcia.margis@ufrgs.br.




Figure 1 - Number of publication evolution addressing antioxidant enzymes in plants in the last ten years. CAT (catalase), SOD (Superoxide dismutase), GPX (Glutathione peroxidase) and APX (Ascorbate peroxidase) enzymes, Reactive Oxygen Species (ROS).

creasing number of publications addressing CAT, SOD, GPX and APX enzymes in plants are examples of this tendency, especially APX, which had the number of articles doubled from 158 published in 2000 to 368 in 2010 (ISI Web of Knowledge database). Publications related more specifically to ROS in plants increased 18 times in the same period. The data presented in this study confirm those reported by Azevedo and Azevedo (2006) in which the number of publications addressing antioxidant mechanisms increased after 2000. This shows the relevance of studying these enzymes to further understand the biological processes dealing with oxidative stress responses in plants. The focus of this review is to discuss the main findings related to the APX enzyme at molecular and physiological levels, in different plant species. *APX* gene modulation in response to abiotic stress conditions, especially temperature, high light, drought, salinity and heavy metals will also be reviewed.

Ascorbate Peroxidase in Plants

Ascorbate peroxidase (APX) (EC 1.11.1.11) belongs to the class I heme-peroxidases that is found in higher plants, chlorophytes (Takeda *et al.*, 1998, 2000), red algae (Sano *et al.*, 2001), and members of the protist kingdom (Shigeoka *et al.*, 1980; Wilkinson *et al.*, 2002). APX and other peroxidase sequences from all kingdoms of life are stored in the database Peroxibase (Oliva *et al.*, 2009), which also provides a series of bioinformatics tools useful for analyzing the peroxidases stored sequences.

Genomic and cDNA *APX* sequences were obtained from a great variety of plant species, showing that APX are widely distributed in the vegetal kingdom. These enzymes are encoded by small gene families in these organisms (Passardi *et al.*, 2007). The different isoforms are classified according to their subcellular localization. Soluble isoforms are found in cytosol (cAPX), mitochondria (mitAPX) and chloroplast stroma (sAPX), while membrane-bound isoforms are found in microbody (including

peroxisome and glyoxisome) (mAPX) and chloroplast thylakoids (tAPX). The presence of organelle-specific targeting peptides and transmembrane domains found in the N- and C- terminal protein regions determine the final subcellular localization of the isoenzyme (Shigeoka *et al.*, 2002; Teixeira *et al.*, 2004, 2006).

Plant chloroplastic APX (chlAPX) isoenzymes encoding genes are divided into two groups. The first group comprises single genes encoding two isoenzymes through a post-transcriptional alternative splicing regulation. This group includes genes from spinach (*S. oleracea*), tobacco (*N. tabacum*), pumpkin (*Cucurbita sp*) and ice plant (*M. crystallium*). In the second group, individual genes codify different isoenzymes which are individually regulated. This group includes genes from *Arabidopsis*, rice, and tomato. The mechanism of alternative splicing in chlAPX has been studied in spinach (Ishikawa and Shigeoka, 2008) and the results showed that alternative splicing is fundamental for controlling the expression of stromal (sAPX) and thylakoid (tAPX) isoenzymes. This regulation occurs in a tissue-dependent manner.

Ascorbate peroxidases have been partially characterized in some plant species. In spinach, the APX family is formed by genes encoding one cytosolic and two chloroplastic (sAPX and tAPX membrane) isoenzymes, one targeted to microbody membrane and an unknown putative cytosol-soluble isoenzyme (Ishikawa *et al.*, 1995, 1996, 1998). In cowpea, four cDNAs were isolated and characterized, corresponding to putative cytosolic, peroxisomal and chloroplastic (thylakoid and stromal) APX isoforms (D'Arcy-Lameta *et al.*, 2006). Six loci encoding *APX* were identified in *Eucalyptus grandis* and their subcellular localizations were indicated by prediction programs. Among the six isoforms, three were putatively identified as cytosolic, one as a putative peroxisomal protein and two predicted to be associated with chloroplasts (Teixeira *et al.*, 2005). In tomato, seven members were identified, three cytosolic, two peroxisomal, and two chloroplastic (Najami *et al.*, 2008). In the model plant *Arabidopsis thaliana*, the presence of nine *APX* genes was described, two chloroplastic, one thylakoid-bound and one member whose product is targeted to both chloroplast stroma and mitochondria (Chew *et al.*, 2003); the intracellular localization of an additional member is yet unknown. In addition, three cytosolic and three microsomal proteins were also described (Mittler *et al.*, 2004; Narendra *et al.*, 2006; Panchuk *et al.*, 2002). In another important model plant, rice, the *APX* gene family comprises eight members, *viz.* two cytosolic, two peroxisomal, two chloroplastic (stromal and thylakoid-bound) and two mitochondrial ones (Teixeira *et al.*, 2004, 2006). Recently, a new protein has also been identified as functionally associated with APX in rice, the APX-R (Ascorbate peroxidase-related) (Lazarotto *et al.*, 2011). Detailed analyses of evolution and structure of *APX-R* genes indi-

cate that these genes correspond to a new class of hemeperoxidases (Lazzarotto *et al.*, 2011).

APX isoenzymes are labile in the absence of AsA. Thus, high level of endogenous AsA is essential to effectively maintain the antioxidant system that protects plants from oxidative damage (Asada, 1992; Shigeoka *et al.*, 2002). Under special conditions in which the concentration of AsA is lower than 20 μ M, the APX activity is quickly lost, this making the chlAPX the least stable isoform. Both cAPX and mAPX have half-inactivation times of around one hour or more, while that for mitAPX and chlAPX is less than 30 seconds (Chen and Asada, 1989; Miyake *et al.*, 1993; Ishikawa *et al.*, 1998; Yoshimura *et al.*, 1998; Leonardis *et al.*, 2000).

APX enzyme Responses Under Abiotic Stress

The expression of *APX* encoding genes is modulated by various environmental stimuli, such as drought and salt stress, high light, high and low temperatures, pathogen attacks, H_2O_2 and abscisic acid (Zhang *et al.*, 1997; Yoshimura *et al.*, 2000; Agrawal *et al.*, 2003; Fryer *et al.*, 2003; Menezes-Benavente *et al.*, 2004; Teixeira *et al.*, 2006; Rosa *et al.*, 2010; Bonifacio *et al.*, 2011). Furthermore, the transcriptional expression of *APX* genes is tissue and developmental stage dependent (Agrawal *et al.*, 2003; Teixeira *et al.*, 2006).

Salt stress

Plants are greatly affected by salinity, which causes alteration in nutrient uptake, accumulation of toxic ions, osmotic stress, and oxidative stress (Verslues *et al.*, 2006). Consequently, salinity results in molecular damage, growth arrest, and even cell death (Wang *et al.*, 2008). Salt stress induces the production of ROS, and the response of *APX* genes to this condition is tissue and developmental stage regulated. When the response of major antioxidant enzymes transcripts was analyzed for different developmental stages in salt stressed rice, *cAPX* was up-regulated in 11-day-old seedlings, while in 6-week-old plants salt had no significant effect on this gene (Menezes-Benavente *et al.*, 2004). In addition to *APX* expression alteration, discrimination on *CAT* transcript accumulation was also noticed in the basal region of rice leaves under salinity (Yamane *et al.*, 2010). Concerning the APX rice isoforms, induction was observed for the *OsAPX1*, *OsAPX4*, *OsAPX6* and *OsAPX7* genes, whereas cytosolic *OsAPX2* gene expression was not altered by salinity (Yamane *et al.*, 2010).

Teixeira *et al.* (2006) reported that three rice *APX* genes, *OsAPX2*, *OsAPX7*, and *OsAPX8*, showed altered transcript levels in response to NaCl treatment. The expression of *OsAPX2* and *OsAPX7* was increased, whereas *OsAPX8* transcript accumulation was strongly suppressed in plants undergoing salt stress (Teixeira *et al.*, 2006). The transcript level of *OsAPX8* was slightly decreased by salin-

ity in the basal region of rice leaves (Yamane *et al.*, 2010). On the other hand, *OsAPX8* expression in rice roots was enhanced by all NaCl concentrations tested (150, 200 and 300 mM), and *OsAPX7* expression was down-regulated by 300 mM NaCl. This discrepancy in regulation for the *OsAPX* genes might be due to differences in cultivars, organs, plant age and growth conditions (Hong *et al.*, 2007). An increase in rice cytosolic *APX2* gene transcript levels after treatment with salt has previously been shown by our group (Menezes-Benavente *et al.*, 2004; Teixeira *et al.*, 2006). In accordance, transgenic *Arabidopsis* plants over-expressing cytosolic *OsAPXb* (*OsAPX2*) showed higher tolerance to NaCl than those over-expressing cytosolic *OsAPXa* (*OsAPX1*) (Lu *et al.*, 2007). A similar increment in salt stress tolerance was also observed in transgenic tobacco over-expressing the *Arabidopsis cAPX* gene (Badawi *et al.*, 2004) and also in tobacco plants over-expressing a *Solanum lycopersicum* thylakoid-bound ascorbate peroxidase gene (*StAPX*) (Sun *et al.*, 2010b). Transgenic tobacco plants that simultaneously expressed CuZnSOD, APX, and DHAR in chloroplasts presented increased protection against salt induced injury (Lee *et al.*, 2007b). Transgenic tobacco BY-2 cells with 50 and 75% lower cAPX activity showed higher intracellular content of ROS. On the other hand, the tobacco cells showed a potential enhancement in tolerance to heat and salt stress, perhaps by induction of stress-related gene expression. However, no substantial differences were observed in the activity levels of the other antioxidant enzymes (Ishikawa *et al.*, 2005).

In barley, the transcript level of peroxisomal *APX* gene (*HvAPX1*) increased significantly under salt stress (Shi *et al.*, 2001). However, *Arabidopsis apx3* knockout mutants exposed to normal or stressful conditions did not present disturbed growth or development. In these plants, other antioxidant enzymes possibly compensate the lack of the peroxisomal isoform (Narendra *et al.*, 2006). In contrast, the overexpression of a *Populus* peroxisomal *APX* (*PpAPX*) gene in transgenic tobacco improved salt tolerance at the vegetative stage and plants were more resistant to oxidative damage induced by methyl viologen (MV) and, in addition, the plants had longer roots (Li *et al.*, 2009). Lin and Pu (2010) studied changes in enzymes involved in ROS scavenging in sweet potato plants tolerant and sensitive to salinity. After exposure to salinity (450 mM NaCl), APX activity increased in plants at 24 and 48 h, and this response was higher in a salt-stress tolerant genotype than in the salt sensitive ones. The expression of *cAPX*, *mAPX* and *chlAPX* in response to salinity was tissue specific and dependent on stress duration (Lin and Pu, 2010). Taken together, these studies put in evidence that salt stress causes disturbances in antioxidant gene expression by producing alterations in the transcriptional pattern in several plant species, and that the expression of distinct APX isoform may result in redox homeostasis regulation in each cellular compartment.

Temperature stress

Extreme temperatures affect the growth, yield and quality of plant production. ROS levels tend to increase if plants are exposed to stressful conditions such as low or high temperatures (Mittler *et al.*, 2004; Scandalios, 2005). In potato tubers, the transient accumulation of *cAPX* mRNA after storage at low-temperature was greater than after high-temperature storage, showing that *APX* expression was induced in response to low temperature (Kawakami *et al.*, 2002). Likewise, the two rice *cAPX* (*OsAPX1* and *OsAPX2*) genes were induced after rice plants were exposed to low temperatures. Furthermore, *OsAPX3*, *OsAPX4*, *OsAPX6* and *OsAPX7* were also significantly induced, while *OsAPX8* were repressed after 24 h under low temperature (unpublished data). The sweet potato *cAPX* gene was highly induced in leaves after exposure to high temperature (Park *et al.*, 2004). In cucumber plants submitted to heat treatment, the activities of *cAPX*, *sAPX* and *mAPX* increased after an initial slight decline during the course of the experiment. The expression of *sAPX* followed a similar pattern (Song *et al.*, 2005). In response to cold, the expression of a peroxisomal *APX* gene increased slightly in *Arabidopsis* (Zhang *et al.*, 1997).

These results were corroborated in *Arabidopsis* by overexpressing a putative peroxisomal membrane-bound *APX* from barley, resulting in an increased tolerance to higher temperature treatment (Shi *et al.*, 2001). Furthermore, the overexpression of chloroplastic *tAPX* in tobacco plants improved the tolerance to chilling stress combined with high light intensity (Yabuta *et al.*, 2002). On the other hand, *Arabidopsis* plants lacking *tAPX* had enhanced tolerance to heat stress (Miller *et al.*, 2007). Recently, Sato *et al.* (2011) showed that transgenic rice plants overexpressing a cytosolic *APX1* gene (*OsAPXa*) which exhibited higher *APX* activity in spikelets than in wild type (WT) plants, sustained higher levels of *APX* activity under cold stress, resulting in enhanced cold tolerance at the booting stage.

Plants with enhanced tolerance to multiple environmental stresses were obtained through induced expression of *CuZnSod* and *APX* genes. *Sod* and *APX* genes were expressed in chloroplasts of potato plants under the control of an oxidative stress inducible promoter - *SWPA2*. These plants showed enhanced tolerance to MV and when exposed to 42 °C for 20 h, the photosynthetic activity of these transgenic plants decreased by only 6%, whereas in non-transformed (NT) plants it decreased by 29% (Tang *et al.*, 2006). Sweet potato plants expressing both *CuZnSod* and *APX* in chloroplasts through the inducible promoter also showed higher tolerance to MV-mediated oxidative stress and chilling stress (Lim *et al.*, 2007). The tolerance to high and low temperature stresses was studied in tobacco plants overexpressing a tomato *tAPX* gene. The overexpression of chloroplastic *APX* played a significant role in H₂O₂ detoxification and in minimizing photooxidative damage during temperature stress. The transgenic plants showed a higher

photochemical efficiency of photosystem II when compared to WT plants under cold and heat stresses (Sun *et al.*, 2010a). These results put in evidence that the manipulation of the antioxidative mechanism in chloroplasts may be applied in the development of plants with increased tolerance to multiple environmental stresses.

High light stress

Plants exposed to excessive light can suffer photo-inhibition, serious damage to the photosynthetic apparatus, and degradation of photosynthetic proteins (Demmig-Adams and Adams III, 1992). Light stress can also lead to ROS accumulation and antioxidant enzymes activation (Mittler, 2002). The responses of *APX* isoenzymes to photooxidative stress were studied in spinach leaves during high light stress. *cAPX* activity and transcripts increased during high light stress, however protein levels were not altered. The activities of *chlAPX* isoforms showed a gradual decrease, while the other isoenzymes showed no significant variation in transcript and protein levels, as well as activities (Yoshimura *et al.*, 2000). In wheat, a mutant line showing decreased *tAPX* activity presented reduced photosynthetic activity and biomass accumulation when growing under high-light intensity, suggesting that *tAPX* is essential for photosynthesis (Danna *et al.*, 2003). Single mutants of *Arabidopsis* lacking *tAPX* or *sAPX* presented higher levels of H₂O₂ and oxidized proteins than WT plants when exposed to high light and MV stresses. The strongest effect of photooxidative stress was observed in plants lacking *tAPX*, these showing increased H₂O₂ accumulation and oxidized proteins (Maruta *et al.*, 2010).

Double mutants deficient in two *APX* genes, thylakoid-bound and a cytosolic one (*tylapx/apx1*), resulted in different signals in *Arabidopsis* plants, such as late flowering, low protein oxidation during light stress and enhanced accumulation of anthocyanins (Miller *et al.*, 2007). Mutants lacking a functional copy of *tAPX*, *sAPX* or both, were characterized in *Arabidopsis* under photooxidative stress during germination. The stress led to chloroplast bleaching in *sapx* single-mutant and *tapx/sapx* double-mutant plants, while the greening process of WT and *tapx* plants was partially impaired (Kangasjarvi *et al.*, 2008). When mature leaves of *tapx/sapx* double mutants were submitted to short-term photooxidative stress induced by high light or MV treatment, the plants showed susceptibility (Kangasjarvi *et al.*, 2008). These results indicate that the *APXs* isoenzymes are indispensable under environmental stresses in different species, especially under light stress conditions.

In *Arabidopsis* leaves, high light treatment induced the expression of cytosolic *APX2*, which has its expression restricted to bundle sheath cells of the vascular tissue (Fryer *et al.*, 2003). In *Arabidopsis*, *APX1* knockout plants showed suppressed growth and development, altered stomatal responses and induction of heat shock proteins

during light stress. The inactivation of cytosolic APX resulted in the alteration of several transcripts involved in different functions. In transgenic APX1 plants kept under optimal conditions, the transcripts encoding APX enzymes were not elevated. However, during light stress, certain enzymes were induced in knockout-APX1 plants (Pnueli *et al.*, 2003). In another study (Davletova *et al.*, 2005) with APX1-deficient *Arabidopsis* plants the observation was that the entire chloroplastic H₂O₂-scavenging system collapsed, H₂O₂ levels increased and protein oxidation occurred in leaves subjected to a moderate light stress, suggesting that the absence of cytosolic APX1 resulted not only in the accumulation of H₂O₂ but also in damage to specific proteins in leaf cells. On the other hand, rice plants double silenced for cytosolic APXs up-regulated other peroxidases, making these transgenic plants able to survive under stress, such as salt, heat, high light and MV, similar to NT plants. The antioxidative compensatory mechanism exhibited by the silenced plants was associated with increased expression of *Gpx* genes. The transcript levels of *OsCatA* and *OsCatB* and the activities of CAT and guaiacol peroxidase (GPOD; type III peroxidases) were also up-regulated. In contrast, none of the other isoforms of *OsAPX* were up-regulated under normal growth conditions. These results suggested that signaling mechanisms triggered in rice could be distinct from those proposed for *Arabidopsis* (Bonifacio *et al.*, 2011).

Drought stress

Drought stress in plants leads to severe effects such as reduction in vegetative growth and cell division. As a consequence of drought stress several changes occur inside the cell, including changes in gene expression levels, synthesis of molecular chaperones, and activation of enzymes involved in the production and removal of ROS (Mahajan and Tuteja, 2005). In two cowpea (*Vigna unguiculata*) cultivars, one drought-tolerant and the other drought-sensitive, APX activity was 60% higher in tolerant plants cultivated under control conditions. In response to drought stress, a higher increase in transcript levels of cytosolic and peroxisomal APX genes was observed in the sensitive cultivar (D'Arcy-Lameta *et al.*, 2006). Chloroplastic APX genes expression was stimulated earlier in the tolerant cultivar when submitted to drought stress. These data suggest the capacity of these enzymes to efficiently detoxify ROS at their production site (D'Arcy-Lameta *et al.*, 2006). Relative APX transcript levels showed distinct changes in two genotypes of wheat exposed to mild water deficit. Cytosolic APX1 expression levels increased in both genotypes, while cytosolic APX2 was up-regulated only in the drought-tolerant genotype. The transcript level of thylakoid APX increased in the drought-tolerant genotype, while stromal APX2 showed higher expression levels in the drought-sensitive cultivar (Secenji *et al.*, 2010).

APX gene expression patterns in rice were studied after 15 days of drought stress. In marked contrast with the experiments with wheat, thylakoid APX (*OsAPX8*) expression was down-regulated in this condition, while the *OsAPX1*, *OsAPX2*, *OsAPX5*, *OsAPX6* and *OsAPX7* genes were up-regulated. The peroxisomal *OsAPX3* gene was not affected, while *OsAPX4* was slightly but significantly down-regulated by this treatment (Rosa *et al.*, 2010). This discrepancy could be due to distinct responses of APX genes in different species and to different magnitudes of stress. In *Arabidopsis*, APX1 protein and mRNA accumulated during combination of heat and drought stress. A cytosolic APX1-deficient mutant accumulated more H₂O₂ and was more sensitive to stress combination than WT plants when exposed to heat and drought stress combined. In contrast, plants deficient in thylakoid APX were not more sensitive to this stress combination than APX1-deficient mutant or WT plants. The cytosolic APX1 gene may thus play a key role in the acclimation of plants to combined stress such drought and heat (Koussevitzky *et al.*, 2008). Indeed, when the overexpression of cytosolic APX was studied in tobacco chloroplasts, its overexpression protected the plant from several oxidative stresses, including drought and polyethylene glycol-induced stress (Badawi *et al.*, 2004). Plants overexpressing other antioxidant enzymes in different species showed increased tolerance to various stresses, including drought resistance. The overexpression of a *Populus* peroxisomal ascorbate peroxidase (*PpAPX*) gene in transgenic tobacco improved drought resistance in these plants (Li *et al.*, 2009). The overexpression of tomato (*Solanum lycopersicum*) thylakoid-bound APX (*StAPX*) gene in tobacco plants enhanced the tolerance of these plants to salt and osmotic stress (Sun *et al.*, 2010b).

Heavy metals

The contamination of soils with heavy metals is a serious environmental problem that limits crop production. Exposure at higher concentrations of heavy metals can increase the production of ROS and change antioxidant response (Gratão *et al.*, 2005). Exposure of pea (*Pisum sativum* L.) plants to cadmium changed enzymatic and non-enzymatic antioxidant defenses, however, APX activity or accumulation of its transcripts were not significantly different (Romero-Puertas *et al.*, 2007). However, it was observed that in coffee cells, the activity of APX was increased at the lower cadmium concentration. On the other hand, APX activity was not detectable in cells submitted to the higher cadmium concentration after 24 h of treatment (Gomes-Junior *et al.*, 2006b).

An increase in APX activity was also observed in response to other heavy metals such as aluminum (Sharma and Dubey, 2007). In rice, the transcript levels of all *OsAPX* genes, except *OsAPX6*, were significantly increased after eight hours of 20 ppm aluminum exposure (Rosa *et al.*, 2010). In pea plants, *cAPX* expression increased in the

shoots under aluminum treatment, but APX activity presented a significant decline at 10 μM aluminum in roots and shoots after 24 and 48 h of stress; at 50 μM aluminum treatment, however, APX activity did not show any significant changes (Panda and Matsumoto, 2010). Transgenic rice plants double silenced for *APX1* and *APX2* (*APX1/2s* plants) exhibited normal development and enhanced tolerance to a toxic concentration of aluminum (Rosa *et al.*, 2010). In bean plants, the expression of cytosolic APX was induced both at mRNA and protein levels in leaves of de-rooted plants in response to iron overload. Likewise, transgenic tobacco plants with suppressed cytosolic APX levels were more sensitive to iron application than WT plants (Pekker *et al.*, 2002). In coffee cells treated with nickel showed a rapid increase in APX activity, although the activity trends were slightly different between the two nickel concentrations tested (0.05 mM and 0.5 mM) (Gomes-Junior *et al.*, 2006a). Transgenic tall fescue plants expressing the CuZnSOD and APX genes in chloroplasts were submitted to copper, cadmium or arsenic treatment. Of the metals tested, copper and cadmium increased SOD

and APX activities in control and transgenic plants, with a higher increase observed in transgenic plants. In contrast, in leaves exposed to arsenic, both enzymes exhibited less activity when compared to other treatments and no significant differences were observed between control and transgenic plants (Lee *et al.*, 2007a). These results emphasize the important role of APX and other antioxidant enzymes in H_2O_2 scavenging under toxic metals levels in the soil.

Conclusions

Ascorbate peroxidase is a key enzyme regulating ROS levels acting in different subcellular compartments (Figure 2). The expression of *APX* encoding genes is differentially modulated by several abiotic stresses in different plant species. All the data collected so far firmly indicate that APX isoforms play important and direct roles as protective elements against adverse environmental conditions. The diverse effects of knockdown or knockout of different *APX* genes on the plant growth, physiology and antioxidant metabolism indicate that APX may also regulate redox signaling pathways involved in plant development. These re-




Figure 2 - APX enzymes and the elimination of ROS excess in different subcellular compartments. H_2O_2 is generated in normal metabolism via the Mehler reaction in chloroplasts, electron transport in mitochondria and photorespiration in peroxisomes. Abiotic and biotic stresses enhance H_2O_2 and chlAPX, mAPX, cAPX and mitAPX enzymes which can eliminate ROS excess in different subcellular compartments. The plasma membrane-NADPH oxidases also generate H_2O_2 , which can cross membranes through aquaporin channels. Superoxide (O_2^-), hydrogen peroxide (H_2O_2), monodehydroascorbate (MDHA).

sults emphasize the importance and complexity of the interactions of APX with other antioxidants in fine tuning the vegetal antioxidant metabolism.

Acknowledgments

The authors are grateful to Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for financial support.

References

- Agrawal GK, Jwa NS, Iwahashi H and Rakwal R (2003) Importance of ascorbate peroxidases OsAPX1 and OsAPX2 in the rice pathogen response pathways and growth and reproduction revealed by their transcriptional profiling. *Gene* 322:93-103.
- Apel K and Hirt H (2004) Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. *Annu Rev Plant Biol* 55:373-399.
- Asada K (1992) Ascorbate peroxidase - A hydrogen peroxide-scavenging enzyme in plants. *Physiol Plantarum* 85:235-241.
- Asada K (1999) The water cycle in chloroplasts: Scavenging of active oxygens and dissipation of excess photons. *Annu Rev Plant Physiol Plant Mol Biol* 50:601-639.
- Azevedo JA and Azevedo RA (2006) Heavy metals and oxidative stress: Where do we go from here? *Commun Biometry Crop Sci* 1:135-138.
- Badawi GH, Kawano N, Yamauchi Y, Shimada E, Sasaki R, Kubo A and Tanaka K (2004) Over-expression of ascorbate peroxidase in tobacco chloroplasts enhances the tolerance to salt stress and water deficit. *Physiol Plant* 121:231-238.
- Bonifacio A, Martins MO, Ribeiro CW, Fontenele AV, Carvalho FE, Margis-Pinheiro M and Silveira JA (2011) Role of peroxidases in the compensation of cytosolic ascorbate peroxidase knockdown in rice plants under abiotic stress. *Plant Cell Environ* 34:1705-1722.
- Chen GX and Asada K (1989) Ascorbate peroxidase in tea leaves: Occurrence of two isozymes and the differences in their enzymatic and molecular properties. *Plant Cell Physiol* 30:897-998.
- Chew O, Whelan J and Millar AH (2003) Molecular definition of the ascorbate-glutathione cycle in Arabidopsis mitochondria reveals dual targeting of antioxidant defenses in plants. *J Biol Chem* 278:46869-46877.
- D'Arcy-Lameta A, Ferrari-Iliou R, Contour-Ansel D, Pham-Thi AT and Zuily-Fodil Y (2006) Isolation and characterization of four ascorbate peroxidase cDNAs responsive to water deficit in cowpea leaves. *Ann Bot* 97:133-140.
- Danna CH, Bartoli CG, Sacco F, Ingala LR, Santa-Maria GE, Guiamet JJ and Ugalde RA (2003) Thylakoid-bound ascorbate peroxidase mutant exhibits impaired electron transport and photosynthetic activity. *Plant Physiol* 132:2116-2125.
- Davletova S, Rizhsky L, Liang H, Shengqiang Z, Oliver DJ, Coutu J, Shulaev V, Schlauch K and Mittler R (2005) Cytosolic ascorbate peroxidase 1 is a central component of the reactive oxygen gene network of Arabidopsis. *Plant Cell* 17:268-281.
- Demmig-Adams B and Adams WW III (1992) Photoprotection and other responses of plants to high light stress. *Annu Rev Plant Physiol and Plant Mol Biol* 43:599-626.
- Foyer CH and Noctor G (2005) Redox homeostasis and antioxidant signaling: A metabolic interface between stress perception and physiological responses. *Plant Cell* 17:1866-1875.
- Fryer MJ, Ball L, Oxborough K, Karpinski S, Mullineaux PM and Baker NR (2003) Control of Ascorbate Peroxidase 2 expression by hydrogen peroxide and leaf water status during excess light stress reveals a functional organisation of Arabidopsis leaves. *Plant J* 33:691-705.
- Gomes-Junior RA, Moldes CA, Delite FS, Gratão PL, Mazzafera P, Lea PJ and Azevedo RA (2006a) Nickel elicits a fast antioxidant response in *Coffea arabica* cells. *Plant Physiol Biochem* 44:420-429.
- Gomes-Junior RA, Moldes CA, Delite FS, Pompeu GB, Gratão PL, Mazzafera P, Lea PJ and Azevedo RA (2006b) Antioxidant metabolism of coffee cell suspension cultures in response to cadmium. *Chemosphere* 65:1330-1337.
- Gratão PL, Polle A, Lea PJ and Azevedo RA (2005) Making the life of heavy metal-stressed plants a little easier. *Funct Plant Biol* 32:481-494.
- Hong CY, Hsu YT, Tsai YC and Kao CH (2007) Expression of ASCORBATE PEROXIDASE 8 in roots of rice (*Oryza sativa* L.) seedlings in response to NaCl. *J Exp Bot* 58:3273-3283.
- Ishikawa T and Shigeoka S (2008) Recent advances in ascorbate biosynthesis and the physiological significance of ascorbate peroxidase in photosynthesizing organisms. *Biosci Biotechnol Biochem* 72:1143-1154.
- Ishikawa T, Sakai K, Takeda T and Shigeoka S (1995) Cloning and expression of cDNA encoding a new type of ascorbate peroxidase from spinach. *FEBS Lett* 367:28-32.
- Ishikawa T, Sakai K, Yoshimura K, Takeda T and Shigeoka S (1996) cDNAs encoding spinach stromal and thylakoid-bound ascorbate peroxidase, differing in the presence or absence of their 3'-coding regions. *FEBS Lett* 384:289-293.
- Ishikawa T, Yoshimura K, Sakai K, Tamoi M, Takeda T and Shigeoka S (1998) Molecular characterization and physiological role of a glyoxysome-bound ascorbate peroxidase from spinach. *Plant Cell Physiol* 39:23-34.
- Ishikawa T, Morimoto Y, Madhusudhan R, Sawa Y, Shibata H, Yabuta Y, Nishizawa A and Shigeoka S (2005) Acclimation to diverse environmental stresses caused by a suppression of cytosolic ascorbate peroxidase in tobacco BY-2 cells. *Plant Cell Physiol* 46:1264-1271.
- Kangasjarvi S, Lepisto A, Hannikainen K, Piippo M, Luomala EM, Aro EM and Rintamaki E (2008) Diverse roles for chloroplast stromal and thylakoid-bound ascorbate peroxidases in plant stress responses. *Biochem J* 412:275-285.
- Kawakami S, Matsumoto Y, Matsunaga A, Mayama S and Mizuno M (2002) Molecular cloning of ascorbate peroxidase in potato tubers and its response during storage at low temperature. *Plant Sci* 163:829-836.
- Koussevitzky S, Suzuki N, Huntington S, Armijo L, Sha W, Cortes D, Shulaev V and Mittler R (2008) Ascorbate peroxidase 1 plays a key role in the response of *Arabidopsis thaliana* to stress combination. *J Biol Chem* 283:34197-34203.
- Lazarotto F, Teixeira FK, Rosa SB, Dunand C, Fernandes CL, de Vasconcelos Fontenele A, Silveira JA, Verli H, Margis R

- and Margis-Pinheiro M (2011) Ascorbate peroxidase-related (APx-R) is a new heme-containing protein functionally associated with ascorbate peroxidase but evolutionarily divergent. *New Phytol* 191:234-250.
- Lee SH, Ahsan N, Lee KW, Kim DH, Lee DG, Kwak SS, Kwon SY, Kim TH and Lee BH (2007a) Simultaneous overexpression of both CuZn superoxide dismutase and ascorbate peroxidase in transgenic tall fescue plants confers increased tolerance to a wide range of abiotic stresses. *J Plant Physiol* 164:1626-1638.
- Lee YP, Kim SH, Bang JW, Lee HS, Kwak SS and Kwon SY (2007b) Enhanced tolerance to oxidative stress in transgenic tobacco plants expressing three antioxidant enzymes in chloroplasts. *Plant Cell Rep* 26:591-598.
- Leonardis S, Dipierro N and Dipierro S (2000) Purification and characterization of an ascorbate peroxidase from potato tuber mitochondria. *Plant Physiol Biochem* 38:773-779.
- Li Y-J, Hai R-L, Du X-H, Jiang X-N and Lu H (2009) Overexpression of a Populus peroxisomal ascorbate peroxidase (PpAPX) gene in tobacco plants enhances stress tolerance. *Plant Breed* 128:404-410.
- Lim S, Kim Y-H, Kim S-H, Kwon S-Y, Lee H-S, Kim J-S, Cho K-Y, Paek K-Y and Kwak S-S (2007) Enhanced tolerance of transgenic sweetpotato plants that express both CuZnSOD and APX in chloroplasts to methyl viologen-mediated oxidative stress and chilling. *Mol Breed* 19:227-239.
- Lin KH and Pu SF (2010) Tissue- and genotype-specific ascorbate peroxidase expression in sweet potato in response to salt stress. *Biol Plantarum* 54:664-670.
- Lu Z, Liu D and Liu S (2007) Two rice cytosolic ascorbate peroxidases differentially improve salt tolerance in transgenic Arabidopsis. *Plant Cell Rep* 26:1909-1917.
- Mahajan S and Tuteja N (2005) Cold, salinity and drought stresses: An overview. *Arch Biochem Biophys* 444:139-158.
- Maruta T, Tanouchi A, Tamoi M, Yabuta Y, Yoshimura K, Ishikawa T and Shigeoka S (2010) Arabidopsis chloroplastic ascorbate peroxidase isoenzymes play a dual role in photoprotection and gene regulation under photooxidative stress. *Plant Cell Physiol* 51:190-200.
- Menezes-Benavente L, Teixeira FK, Kamei CLA and Margis-Pinheiro M (2004) Salt stress induces altered expression of genes encoding antioxidant enzymes in seedlings of a Brazilian indica rice (*Oryza sativa* L.). *Plant Sci* 166:323-331.
- Miller G, Suzuki N, Rizhsky L, Hegie A, Koussevitzky S and Mittler R (2007) Double mutants deficient in cytosolic and thylakoid ascorbate peroxidase reveal a complex mode of interaction between reactive oxygen species, plant development, and response to abiotic stresses. *Plant Physiol* 144:1777-1785.
- Mittler R (2002) Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci* 7:405-410.
- Mittler R, Vanderauwera S, Gollery M and Van Breusegem F (2004) Reactive oxygen gene network of plants. *Trends Plant Sci* 9:490-498.
- Miyake C, Cao WH and Asada K (1993) Purification and molecular properties of thylakoid-bound ascorbate peroxidase in spinach chloroplasts. *Plant Cell Physiol* 34:881-889.
- Najami N, Janda T, Barriah W, Kayam G, Tal M, Guy M and Volokita M (2008) Ascorbate peroxidase gene family in tomato: Its identification and characterization. *Mol Genet Genomics* 279:171-182.
- Narendra S, Venkataramani S, Shen G, Wang J, Pasapula V, Lin Y, Korniyev D, Holaday AS and Zhang H (2006) The Arabidopsis ascorbate peroxidase 3 is a peroxisomal membrane-bound antioxidant enzyme and is dispensable for Arabidopsis growth and development. *J Exp Bot* 57:3033-3042.
- Noctor G and Foyer CH (1998) Ascorbate glutathione: Keeping active oxygen under control. *Annu Rev Plant Physiol Plant Mol Biol* 49:249-279.
- Oliva M, Theiler G, Zamocky M, Koua D, Margis-Pinheiro M, Passardi F and Dunand C (2009) PeroxiBase: A powerful tool to collect and analyse peroxidase sequences from Viridiplantae. *J Exp Bot* 60:453-459.
- Panchuk II, Volkov RA and Schöffl F (2002) Heat stress- and heat shock transcription factor-dependent expression and activity of ascorbate peroxidase in Arabidopsis. *Plant Physiol* 129:838-853.
- Panda SK and Matsumoto H (2010) Changes in antioxidant gene expression and induction of oxidative stress in pea (*Pisum sativum* L.) under Al stress. *Biometals* 23:753-762.
- Park SY, Ryu SH, Jang IC, Kwon SY, Kim JG and Kwak SS (2004) Molecular cloning of a cytosolic ascorbate peroxidase cDNA from cell cultures of sweet potato and its expression in response to stress. *Mol Genet Genomics* 271:339-346.
- Passardi F, Bakalovic N, Teixeira FK, Margis-Pinheiro M, Penel C and Dunand C (2007) Prokaryotic origins of the non-animal peroxidase superfamily and organelle-mediated transmission to eukaryotes. *Genomics* 89:567-579.
- Pekker I, Tel-Or E and Mittler R (2002) Reactive oxygen intermediates and glutathione regulate the expression of cytosolic ascorbate peroxidase during iron-mediated oxidative stress in bean. *Plant Mol Biol* 49:429-438.
- Pnueli L, Liang H, Rozenberg M and Mittler R (2003) Growth suppression, altered stomatal responses, and augmented induction of heat shock proteins in cytosolic ascorbate peroxidase (Apx1)-deficient Arabidopsis plants. *Plant J* 34:187-203.
- Romero-Puertas MC, Corpas FJ, Rodríguez-Serrano M, Gómez M, Río LAd and Sandalio LM (2007) Differential expression and regulation of antioxidative enzymes by cadmium in pea plants. *J Plant Physiol* 164:1346-1357.
- Rosa SB, Caverzan A, Teixeira FK, Lazzarotto F, Silveira JA, Ferreira-Silva SL, Abreu-Neto J, Margis R and Margis-Pinheiro M (2010) Cytosolic APx knockdown indicates an ambiguous redox responses in rice. *Phytochemistry* 71:548-558.
- Sano S, Ueda M, Kitajima S, Takeda T, Shigeoka S, Kurano N, Miyachi S, Miyake C and Yokota A (2001) Characterization of ascorbate peroxidases from unicellular red alga *Galdieria partita*. *Plant Cell Physiol* 42:433-440.
- Sato Y, Masuta Y, Saito K, Murayama S and Ozawa K (2011) Enhanced chilling tolerance at the booting stage in rice by transgenic overexpression of the ascorbate peroxidase gene, OsAPXa. *Plant Cell Rep* 30:299-406.
- Scandalios JG (2005) Oxidative stress: Molecular perception and transduction of signals triggering antioxidant gene defenses. *Braz J Med Biol Res* 38:995-1014.
- Secenji M, Hideg E, Bebes A and Gyorgyey J (2010) Transcriptional differences in gene families of the ascorbate-gluta-

- thione cycle in wheat during mild water deficit. *Plant Cell Rep* 29:37-50.
- Sharma P and Dubey RS (2007) Involvement of oxidative stress and role of antioxidative defense system in growing rice seedlings exposed to toxic concentrations of aluminum. *Plant Cell Rep* 26:2027-2038.
- Shi WM, Muramoto Y, Ueda A and Takabe T (2001) Cloning of peroxisomal ascorbate peroxidase gene from barley and enhanced thermotolerance by overexpressing in *Arabidopsis thaliana*. *Gene* 273:23-27.
- Shigeoka S, Ishikawa T, Tamoi M, Miyagawa Y, Takeda T, Yabuta Y and Yoshimura K (2002) Regulation and function of ascorbate peroxidase isoenzymes. *J Exp Bot* 53:1305-1319.
- Shigeoka S, Nakano Y and Kitaoka S (1980) Metabolism of hydrogen peroxide in *Euglena gracilis* Z by L-ascorbic acid peroxidase. *Biochem J* 186:377-380.
- Song XS, Hu WH, Mao WH, Ogwenjo JO, Zhou YH and Yu JQ (2005) Response of ascorbate peroxidase isoenzymes and ascorbate regeneration system to abiotic stresses in *Cucumis sativus* L. *Plant Physiol Biochem* 43:1082-1088.
- Sun W-H, Duan M, Li F, Shu D-F, Yang S and Meng Q-W (2010a) Overexpression of tomato tAPX gene in tobacco improves tolerance to high or low temperature stress. *Biol Plantarum* 54:614-620.
- Sun WH, Duan M, Shu DF, Yang S and Meng QW (2010b) Over-expression of StAPX in tobacco improves seed germination and increases early seedling tolerance to salinity and osmotic stresses. *Plant Cell Rep* 29:917-926.
- Takeda T, Yoshimura K, Ishikawa T and Shigeoka S (1998) Purification and characterization of ascorbate peroxidase in *Chlorella vulgaris*. *Biochimie* 80:295-301.
- Takeda T, Yoshimura K, Yoshii M, Kanahoshi H, Miyasaka H and Shigeoka S (2000) Molecular characterization and physiological role of ascorbate peroxidase from halotolerant *Chlamydomonas* sp. W80 strain. *Arch Biochem Biophys* 376:82-90.
- Tang L, Kwon SY, Kim SH, Kim JS, Choi JS, Cho KY, Sung CK, Kwak SS and Lee HS (2006) Enhanced tolerance of transgenic potato plants expressing both superoxide dismutase and ascorbate peroxidase in chloroplasts against oxidative stress and high temperature. *Plant Cell Rep* 25:1380-1386.
- Teixeira FK, Menezes-Benavente L, Margis R and Margis-Pinheiro M (2004) Analysis of the molecular evolutionary history of the ascorbate peroxidase gene family: Inferences from the rice genome. *J Mol Evol* 59:761-770.
- Teixeira FK, Menezes-Benavente L, Galvão VC and Margis-Pinheiro M (2005) Multigene families encode the major enzymes of antioxidant metabolism in *Eucalyptus grandis* L. *Genet Mol Biol* 28:529-538.
- Teixeira FK, Menezes-Benavente L, Galvão VC, Margis R and Margis-Pinheiro M (2006) Rice ascorbate peroxidase gene family encodes functionally diverse isoforms localized in different subcellular compartments. *Planta* 224:300-314.
- Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu J and Zhu JK (2006) Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *Plant J* 45:523-539.
- Vranova E, Inze D and Van Breusegem F (2002) Signal transduction during oxidative stress. *J Exp Bot* 53:1227-1236.
- Wang R, Chen S, Zhou X, Shen X, Deng L, Zhu H, Shao J, Shi Y, Dai S, Fritz E, Huttermann A and Polle A (2008) Ionic homeostasis and reactive oxygen species control in leaves and xylem sap of two poplars subjected to NaCl stress. *Tree Physiol* 28:947-957.
- Wilkinson SR, Obado SO, Mauricio IL and Kelly JM (2002) *Trypanosoma cruzi* expresses a plant-like ascorbate-dependent hemoperoxidase localized to the endoplasmic reticulum. *Proc Natl Acad Sci USA* 99:13453-13458.
- Yabuta Y, Motoki T, Yoshimura K, Takeda T, Ishikawa T and Shigeoka S (2002) Thylakoid membrane-bound ascorbate peroxidase is a limiting factor of antioxidative systems under photo-oxidative stress. *Plant J* 32:915-925.
- Yamane K, Mitsuya S, Taniguchi M and Miyake H (2010) Transcription profiles of genes encoding catalase and ascorbate peroxidase in rice leaf tissues under salinity. *Plant Prod Sci* 13:164-168.
- Yoshimura K, Ishikawa T, Nakamura Y, Tamoi M, Takeda T, Tada T, Nishimura K and Shigeoka S (1998) Comparative study on recombinant chloroplastic and cytosolic ascorbate peroxidase isozymes of spinach. *Arch Biochem Biophys* 353:55-63.
- Yoshimura K, Yabuta Y, Ishikawa T and Shigeoka S (2000) Expression of spinach ascorbate peroxidase isoenzymes in response to oxidative stresses. *Plant Physiol* 123:223-234.
- Zhang H, Wang J, Nickel U, Allen RD and Goodman HM (1997) Cloning and expression of an Arabidopsis gene encoding a putative peroxisomal ascorbate peroxidase. *Plant Mol Biol* 34:967-971.

Internet Resources

Oliva *et al.*, (2009) Peroxibase, <http://peroxibase.isb-sib.ch> (May 30, 2011).

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.